

Materials for a Flora of the Malayan Peninsula.—By GEORGE KING, M. B., LL. D., F.R.S., C.I.E., *Superintendent of the Royal Botanic Garden, Calcutta.*

[Read June 7th].

No. 5.

ORDER XVI. DIPTEROCARPEÆ.

Resinous trees, rarely climbing shrubs. *Leaves* alternate, simple, quite entire, rarely sinuate-crenate, penni-nerved, the main nerves bold; stipules usually small and inconspicuous, sometimes larger and persistent, or fugitive, leaving an annular scar, (absent in *Ancistrocladus*). *Flowers* in few- or many-flowered, axillary and terminal racemes or panicles. *Bracts* usually minute or 0, rarely larger and persistent. *Sepals* free, or cohering into a tube surrounding but free from, or more or less adnate to, the base of the ovary and fruit. *Petals* contorted, connate at the base, or free. *Stamens* ∞ , 15, 10 or 5, hypogynous or sub-perigynous, free, connate, or adnate to the petals; filaments short, often dilated at the base; anthers 2-celled, the outer valves sometimes larger, connective often aristate or with an obtuse appendage. *Ovary* slightly immersed in the torus, usually 3- rarely 2- or 1-celled; style subulate or fleshy, entire or with 3 minute stigmatic lobes; ovules anatropous, 2 in each cell, pendulous or laterally affixed (solitary and erect in *Ancistrocladus*). *Fruit* usually nut-like, its pericarp leathery or woody, 1- rarely 2-seeded, surrounded by the variously accrescent calyx of which two or more sepals or lobes are usually developed into linear wings. *Seed* exalbuminous (albumen fleshy and ruminant in *Ancistrocladus*); cotyledons fleshy, equal or unequal, straight or more or less plaited and crumpled, sometimes lobed; radicle directed towards the hilum, usually included between the cotyledons.—DISTRIB. Confined (except a few Tropical African species) to Tropical Eastern Asia; genera about 18, species about 250.

Sect. I. EU-DIPTEROCARPEÆ. *Ovaries* 3-celled, each cell 2-ovuled: stigmas united, more or less 3-lobed: seeds usually exalbuminous the outer segments of the fruiting calyx usually enlarged: trees or erect shrubs, mostly stipulate.

Fruiting calyx with 2 or more of its segments or sepals produced into long membranous, reticulate, nerved wings much longer than the fruit; pericarp leathery, (woody in some sp. of *Shorea*).

- Fruiting calyx with a distinct tube.
 Calyx-tube quite free from the fruit 1 *Dipterocarpus*.
 Calyx-tube adherent to the fruit 2 *Anisoptera*.
- Sepals united at the base only, the short calyx-tube either quite free from the fruit or slightly adherent to it, the calyx-segments or sepals valvate or nearly so.
 Stamens with a single, long apical, appendage from the connective 3 *Vatica*.
 Stamens with 4 apical appendages from the anthers and 1 from the connective ... 4 *Pentacme*.
- Sepals free, imbricate.
 The three outer sepals always, and one or both of the inner two occasionally, winged in the fruit; anthers with a short apical appendage from the connective 5 *Shorea*.
 The two outer sepals winged in the fruit, the three inner not longer than the fruit and closely embracing it; stamens with a terminal appendage from the connective longer than the anther 6 *Hopea*.
- Sepals of fruiting-calyx all enlarged but not exceeding, or only slightly exceeding, the fruit; pericarp leathery or woody.
 Fruiting calyx embracing the fruit but not adherent to it.
 Sepals of fruiting calyx slightly thickened.
 Sepals of fruiting-calyx oblong, nearly equal, usually shorter than the fruit, reflexed or erect ... 7 *Retinodendron*.
 Sepals of fruiting-calyx rotund, unequal (the inner two smaller), reflexed ... 8 *Isoptera*.

Sepals of fruiting calyx much thickened and woody at the base.

Calyx forming a cup at the base of the fruit, but not adhering to it: pericarp woody 9 *Balanocarpus*.

Calyx adherent to the fruit: pericarp thickly leathery 10 *Pachynocarpus*.

Sect. II. ANCISTROCLADEÆ. *Ovary* 1-celled with a single ovule; stigmas 3, distinct: *Seeds* with copious ruminant albumen. Exstipulate climbers. 11 *Ancistrocladus*.

1. DIPTEROCARPUS, Gærtn. f.

Lofty trees, stellately pubescent or more or less clothed with fasciated hairs. *Leaves* coriaceous, entire or sinuate-crenate; lateral nerves connected by marginal loops and transverse reticulations; stipules large, valvate, enclosing the terminal bud, finally caducous and leaving an annular scar. *Flowers* large, white or reddish. *Calyx-tube* free. *Petals* usually pubescent externally, especially on the outer margin. *Stamens* ∞; anthers linear, equivalved, acuminate. *Ovary* 3-celled; style filiform; ovules 2 in each cell. *Fruit* nut-like, 1-seeded, enclosed in the accrescent calyx-tube, free; accrescent calyx-lobes 2, erect. *Seed* adnate to the base of the pericarp; cotyledons large, thick, unequal; radicle inconspicuous.—DISTRIB. Tropical E. Asia; species about 60.

Ripe fruit spheroidal or ellipsoidal, neither angled nor winged.

Young branches, petioles, under surfaces of the midribs, and nerves of the leaves covered with coarse stiff fasciculate hairs.

- Fruit glabrous 1. *D. crinitus*.
- „ stellate-pubescent 2. *D. Scortechinii*.

Young branches deciduously pubescent.

Leaves with 12 or more pairs of nerves.

Leaves oblong-elliptic, their under surfaces sparsely stellate-pubescent 3. *D. Skinneri*.

Leaves elliptic or ovate-elliptic, their under surfaces puberulous or quite glabrous 4. *D. turbinatus*.

Leaves with 8 to 10 pairs of nerves.

All parts quite glabrous 5. *D. Kerrii*.

- Ripe fruit with 5 angular tuberosities on its upper portion 6. *D. cornutus*.
- Ripe fruit 5-angled :
- Calyx-tube glabrous ; leaves 2.5 to 3.25 in. long 7. *D. fagineus*.
- Calyx-tube densely stellate-tomentose ; leaves 6 to 8 in. long 8. *D. oblongifolius*.
- Ripe fruit with its 5 angles produced into wings :
- Leaves glabrous :
- Young branches at first scurfy-puberulous, ultimately quite glabrous : buds ovoid, minutely pale canescent 9. *D. grandiflorus*.
- Young branches as in the last, but with conspicuous tawny-tomentose, oblique annuli ; buds cylindric, hoary-canescenscent 10. *D. Kunstleri*.
- Young branches minutely tawny-pubescent, not annulated and never glabrous ; buds ovoid, densely sericeous 11. *D. Griffithii*.
- Leaves minutely stellate-pubescent on the lower surface :
- Flowers about 1 in. long ; leaves with rounded or sub-cordate bases ; young branches very stout, with ovoid buds : the accrescent lobes of the calyx 1.5 in. broad 12. *D. incanus*.
- Flowers 1.5 in. long ; leaves with rounded or cuneate, not sub-cordate, bases : young branches moderately stout with cylindric buds : accrescent calyx-lobes .7 to .8 in. broad 13. *D. alatus*.

1. DIPTEROCARPUS CRINITUS, Dyer in Hook. fil. Fl. Br. Ind. I. 296. A tree 90 to 150 feet high : young branches, petioles, under surface of midrib and nerves, pedicels and outer surface of bracts of inflorescence clothed with stiff yellowish-brown fascicled hairs. *Leaves* very coriaceous, ovate or more usually obovate, acute, the base rounded or subacute ; the edge entire, fringed with fascicled hairs, recurved (at least when dry) ; both surfaces sparsely hispid when young, glabrescent when old ; main nerves 12 to 18 pairs, spreading, rather straight, very prominent on the lower, depressed on the upper, surface ; length 3 to 5 in., breadth 1.75 to 2.75 in., petiole 1 to 1.25 in. *Racemes* about 6-flowered. *Flowers* nearly 2 in. long. *Calyx* glaucous, glabrous. *Petals*

puberulous, linear, blunt. *Stamens* 15. *Fruit* (immature) ellipsoid, wingless, glaucous, smooth; the enlarged calyx-lobes linear-oblong, blunt, 3-nerved, inconspicuously reticulate, shining, 3·5 in. long and ·6 to ·8 in. broad. Dyer in Journ. Bot. 1874, p. 103. *D. hirtus*, Vesque, Comptes-Rendus, 1874, 78, p. 627; Journ. Bot. 1874, p. 151; Dyer l. c. 154.

Malacca; Maingay (Kew Distrib.) No. 196.

Perak; Scortechini, No. 1955. DISTRIB. Borneo: (fide Dyer), Beccari, 779, 1883.

Burck (Ann. Jard. Bot. Buitenzorg, Vol. 6, p. 196) reduces this to *D. Tamparan*, Korth. Korthals however describes the fruit of that species as having accrescent calyx-lobes 13 inches long by 3 broad.

2. *DIPTEROCARPUS SCORTECHINII*, King, n. sp. A large tree: young branches rather stout, densely clothed, (as are the short cylindric buds, the petioles and racemes) with large tufts of coarse, brownish, shining hairs. *Leaves* coriaceous, elliptic-ovate, or sometimes elliptic-sub-ovate, sub-entire, abruptly and shortly acuminate, slightly narrowed to the rounded base; upper surface glabrous or glabrescent, the nerves sparsely stellate-pubescent, the midrib tomentose; under surface sparsely stellate-pubescent, the nerves (and especially the midrib) with long silky hairs intermixed: main nerves 16 to 18 pairs, straight, oblique, very prominent beneath: length 6 to 7·5 in., breadth 3 to 3·5 in., petiole 1 to 1·2 in. *Racemes* few-flowered, short. *Fruit* (? immature) ovoid, contracted under the mouth, glaucous, stellate-pubescent, ·75 in. long and ·5 in. in diam; accrescent calyx-lobes linear-oblong, reticulate, slightly narrowed in the lower half, the apex obtuse, obscurely 3-nerved (the middle nerve bold, the two lateral faint), 4 to 5 in. long and ·8 to 1 in. broad.

Perak; Scortechini, No. 1813.

This is closely allied to *D. crinitus*, Dyer, to which Scortechini doubtfully referred it. It differs from *D. crinitus* in its larger leaves and stellate-pubescent fruit. It has also a different time of flowering; for, as Scortechini remarks in his field notes, this is in immature fruit in the beginning of March, while *D. crinitus* does not come into flower until the end of April.

3. *DIPTEROCARPUS SKINNERI*, King, n. sp. A tall tree; young branches thin, deciduously tawny-pubescent. *Buds* cylindric, narrow, golden-sericeous. *Leaves* oblong-elliptic, narrowed in the upper half or third to the acute or shortly acuminate apex, slightly narrowed to the rounded base, upper surface glabrous or sparsely adpressed-pubescent, the midrib tomentose, the lower sparsely stellate-pubescent, the midrib and 16 to 19 pairs of straight oblique nerves adpressed-sericeous; nerves prominent on the lower, faint on the upper, surface when dry:

length 5 to 8 in., breadth 2·25 to 3 in.; petiole ·7 to ·9 in., tomentose. *Racemes* simple, short, 2- or 3-flowered, pubescent. *Flowers* 2·5 in. long. *Calyx* with narrowly campanulate tube, covered outside with minute, pale, stellate tomentum. *Petals* linear-oblong, blunt, more or less pubescent outside. *Fruit* (? immature) globular-ovoid, glabrous, ·65 in. in diam.: accrescent calyx-lobes glabrous, reticulate, linear, blunt, contracted at the very base, nearly 5 in. long and about ·75 in. broad.

Penang; at the back of West Hill, at an elevation of 1,000 feet. Curtis No. 1403.

A very distinct species known only by Mr. Curtis' scanty specimens. I have named it in honour of Mr. Skinner, Resident Councillor of Penang.

4. *DIPTEROCARPUS TURBINATUS*, Gaertn. f. *Fruct.* III. 51, t. 188. A tree 80 to 100 feet high: young shoots rather slender, at first minutely velvety, pale grey, afterwards glabrous: buds cylindric, softly pale pubescent. *Leaves* thinly coriaceous, elliptic or ovate-elliptic, acute or shortly acuminate, the base rounded or sub-cordate, the edges slightly undulate, sometimes sub-crenate; both surfaces glabrous, or the lower puberulous especially on the midrib and nerves: main nerves 12 to 18 pairs, straight, oblique, prominent on the lower surface; length 4·5 to 11 in., breadth 2·5 to 5·25 in.; petiole 1 to 1·5 in., glabrous or pubescent: stipules tawny-velvety in the lower part but pubescent towards the apex. *Racemes* 3- to 5-flowered. *Flowers* 1·25 to 1·5 in. long. *Calyx*-tube obconic, glabrous, smooth, not winged. *Petals* linear-oblong, obtuse, more or less canescent. *Fruit* ellipsoid-ovoid, tapering to each end when young: globular when ripe and ·75 in. in diam., with neither wings nor ridges; the two accrescent calyx-lobes glabrous, conspicuously reticulate, obscurely 3-nerved, oblong-lanceolate, obtuse, 4 to 4·5 in. long and 1·25 in. broad; the three small lobes of the calyx deltoid, very short. Roxb. *Hort. Beng.* 42; *Fl. Ind.* II. 612; *Corom. Plants* III. 10 t. 213. Ham. in *Mem. Wern. Soc.* VI. 300: *Wall Cat.* 952; A. DC. *Prod.* XVI. 2, 607; W. and Arn. *Prod.* 85; Dyer in *Hook. fil. Fl. Br. Ind.* I, 295: *Journ. Bot.* 1874, p. 102 t. 143, fig. 13: *Kurz. For. Fl. Burm.* I. 114. *D. laevis*, Ham. l. c. 299.; A. DC. l. c. 607. W. and A. *Prod.* 85: *Kurz, l. c.* 114. ?*D. indicus*, Bedd. *Forest. Rep.* 1864-5, 17 cum tab.; *Flora Sylvat.* t. 94.

Assam, Cachar, Chittagong, Burmah, S. India.

VAR. andamanica: enlarged calyx-lobes linear-oblong, not oblanceolate, ·75 in. broad; leaves broadly ovate, sub-cuneate at the base.

South Andaman: common.

Following Dyer, I have included under this the plant named *D. laevis* by Buchanan Hamilton in the *Memoirs of the Wernerian Society*,

Vol. VI. p. 299. Hamilton distinguishes his species *D. laevis* by its flattened branchlets, and perfectly glabrous leaves and petioles, while *D. tuberculatus* Gaertn. has terete branches and pubescent leaves and petioles. The former (called *Dulia Garjan*, by the natives of Chittagong) yields, he says, no wood-oil; while the latter (called *Telia Garjan*) does. The materials before me do not enable me to differentiate the two as species. Moreover, specimens sent to me by Dr. E. Thurston, Reporter on Economic Products to the Government of India, (and which had been collected by the Forest Officer of Chittagong under the vernacular names *Dulia* and *Telia Garjan*) appear exactly alike. Careful investigation in the field may however prove that there is some better basis for Hamilton's view than the trifling differences which he has noted in the outline of the branchlets and the pubescence of the leaves. I am not at all satisfied that the Southern Indian tree named *D. indicus* by Beddome is rightly reduced here. Better Herbarium specimens than any which I have seen, and investigation in the field, are I think required to settle this point also.

5. DIPTEROCARPUS KERRII, King, n. sp. A tall tree; all parts, except the petals, glabrous; young branches thin, slightly flattened at the tips, not annular. *Buds* narrow, cylindric. *Leaves* coriaceous, ovate-elliptic, acute or very shortly and bluntly acuminate, the edges undulate, the base cuneate; main nerves 8 to 11 pairs, oblique, straight, bold and shining on the lower surface; length 3 to 4 in., breadth 2 to 2.5 in., petiole .9 to 1.1 in. *Panicles* short, spreading, few-flowered. *Flowers* 1.5 in. long. *Calyx-tube* glaucous. *Petals* linear-oblong, obtuse, more or less pubescent or tomentose towards their middle externally. *Fruit* turbinate, smooth, 1 to 1.15 in. in diam.; accrescent calyx-lobes linear-oblong, blunt, reticulate, 3-nerved, 4.5 to 5 in. long, and 1.25 to 1.5 in. broad: minor lobes very short, broad, rounded.

Malacca; Maingay (Kew Distrib.) No. 199, Griffith 727, Derry 1032. Pangkore; on Gunong Yunggal, Curtis No. 1561.

Mr. Curtis describes this as a very large tree yielding an oil. It resembles *D. Hasseltii*, Bl., but has much smaller leaves.

I have named this species in honour of Dr. Kerr, an enthusiastic Botanist much interested in the Malayan Flora. Closely allied to this, and perhaps identical with it, is the tree represented by Mr. Curtis' specimen (Waterfall, Penang) No. 1653. The young wood of the latter is however paler than that of *D. Kerrii* from Pangkore and Malacca, and the leaves are puberulous, not glabrous, beneath. I have seen no flowers of it.

6. DIPTEROCARPUS CORNUTUS, Dyer in Hook. fil. Fl. Br. Ind. I, 296. A tree 50 to 70 feet high: young branches stout, compressed, minutely

rufous-tomentose with a few scattered longer hairs. *Leaves* large, coriaceous, oblong, blunt at each end, the edges undulate or obscurely sinuate-crenate: upper surface glabrous, the midrib and nerves pale when dry: under surface densely covered with minute, pale, stellate tomentum: main nerves 16 to 20 pairs, prominent, spreading, straight, the transverse veins rather distinct: length 9 to 14 in., breadth 5 to 8 in., petiole 2 to 3 in.; stipules rufous-sericeous, the hairs fascicled. *Racemes* 7- or 8-flowered. *Flowers* 1·75 in long. *Calyx-tube* 5-winged, canescent, the short lobes very obtuse. *Petals* oblong or sub-spathulate, stellate-canescant. *Fruit* about 1 in. long, sub-globular, with 5 thick short wings in its upper half; enlarged calyx-lobes linear, obtuse, 5 or 6 in. long and 1·25 to 1·75 in. broad, shining, boldly 3-nerved, reticulate. Dyer in Journ. Bot. 1874, p. 103, t. 143, fig. 15. *Parinarium dillenifolium*, R. Br. Wall. Cat. No. 7520. *Petrocarya dillenifolia*, Steud. Nomencl. II, 309.

Singapore: Wallich. Malacca: Maingay (Kew Distrib.) No. 197. Penang: Curtis No. 1402. Perak: Wray, No. 4160.

It was Sir Joseph Hooker who first pointed out that the Wallichian plant No. 7520, issued as *Parinarium*, belongs really to this species.

7. DIPTEROCARPUS FAGINEUS, Vesque in Comptes-Rendus, tome 78, p. 626: Journ. Bot. for 1874, p. 149. A tree 40 to 80 feet high: young branches slender, at first minutely pulverulent tawny-pubescent, ultimately glabrescent or glabrous and dark-coloured, the buds cylindrical. *Leaves* coriaceous, elliptic-ovate to elliptic-lanceolate, acute, the edges entire or sub-undulate-crenulate, the base cuncate, both surfaces puberulous especially on the midrib and nerves; main nerves 10 to 13 pairs, straight, oblique, prominent on the sub-glaucous lower surface; length 2·5 to 3·25 in., breadth 1·3 to 1·75. *Racemes* slender, 1- to 4-flowered. *Flowers* about 1·25 in. long. *Calyx-tube* campanulate, not constricted at the mouth, 5-angled. *Ripe fruit* ellipsoid, tapering more at the base than at the apex, 5-angled, glaucous, 1 in. long: accrescent calyx-lobes linear-oblong, obtuse, contracted at the base, 3-nerved, 2·5 to 3 in. long and about ·75 in. broad. *D. prismaticus*, Dyer Journ. Bot. 1874, pp. 104, 152, t. 144 fig. 17. *Dipterocarpus*, sp. Hook. fil. in Linn. Trans. XXIII, 161.

Perak: King's Collector No. 3527, Scortechini. Penang; Curtis No. 1401.

D. fagineus, Vesque, has been collected hitherto only in Borneo (Beccari No. 3008 and Motley No. 143,) and the leaves are described by Dyer as being papyraceous in texture and having about 8 pairs of lateral nerves. The leaves of the Perak tree which I now refer to this

species, are coriaceous and have 10 to 13 pairs of nerves. The Perak plant may therefore belong to a distinct, but closely allied, species. Curtis' Penang specimens (No. 1401) are quite glabrous in all parts except the petals.

8. *DIPTEROCARPUS OBLONGIFOLIUS*, Blume, Mus. Bot. Lugd. Bat. II, 36. A tall tree: young branches glabrous, dark-coloured, sparsely lenticellate; buds cylindric. *Leaves* coriaceous, oblong or elliptic-oblong, shortly and bluntly acuminate, the edges sub-undulate, the base cuneate; both surfaces shining, glabrous, the midrib and 13 to 16 pairs of straight bold nerves with a few stellate hairs along their sides: length 6 to 8 in., breadth 2 to 2.75 in., petiole .9 to 1.1 in. *Racemes* slightly supra-axillary, densely tawny-tomentose, bifurcating, each branch with 3 to 5 flowers and several linear-membranous deciduous bracts. *Flowers* about 2.5 in. long. *Calyx-tube* fusiform, slightly contracted at the mouth, 1 in. long, boldly 5-angled, densely stellate tawny-tomentose as are the 3 minor calyx lobes; the 2 larger linear-ob lanceolate lobes sparsely stellate-pubescent, boldly 1-nerved and with 2 obscure lateral nerves. *Ripe fruit* unknown. Miq., Fl. Ind. Bat. I, pt. 2, p. 498; A. DC. Prod. XXI. 2, 614; Dyer in Journ. Bot. 1874, 105. *D. stenopterus*, Vesque, Comptes-Rendus, tome 78, p. 625; Journ. Bot. 1874, p. 150.

Perak, Scortechini. **DISTRIB.** Borneo, Sumatra.

Except as regards inflorescence, the Perak specimens of this are practically glabrous. In Bornean specimens, however, the young parts, buds and petioles are fusco-tomentose. (Dyer l. c.)

9. *DIPTEROCARPUS GRANDIFLORUS*, Blanco, Fl. Philipp. Ed. 2, 314. A tree 80 to 120 feet high: young branches rather stout, sub-compressed, at first hoary-puberulous, but finally quite glabrous, nearly black when dry; leaf-buds shortly ovoid, minutely pale-canescens. *Leaves* coriaceous, ovate-elliptic, shortly acuminate; the base broad, rounded or sub-truncate, sub-cordate; the edges entire or obscurely undulate-crenate, both surfaces glabrous; main nerves 14 to 16 pairs, spreading, rather straight, prominent on the lower, obsolete on the upper, surface; length 6 to 9 in., breadth 3.5 to 5 in.; petiole 2 to 3 in. long, glabrous. *Racemes* about 4-flowered. *Flowers* articulated to the rachis, 2 in. long. *Calyx-tube* 5-winged from base to apex. *Petals* linear-oblong. *Fruit* oblong, 2.5 in. long, wings stout, .5 in. or more in width; the 2 accrescent lobes of the calyx oblong, obtuse, glabrous, reticulate, 3-nerved, the mesial nerve the longest and most distinct, 7 to 9 in. long and 1.5 to 2 in. broad, the smaller calyx lobes sub-orbicular. A. DC. Prod. XVI., 2 p. 612; Dyer in Journ. Bot. 1874, p. 106, t. 145, fig 19; Burek in Ann. du Jard. Bot. Buitenzorg, vol 6, 201. *D. Blancoi*, Bl., Mus. Lugd. Bat. II.

35. *D. Motleyanus*, Hook. fil. in Trans. Linn. Soc. XXIII. 159. A.D.C. in DC. Prod. XVI., pt. 2, 611. *D. pterygocalyx*, Scheff. Obs. Phyt. II. 35; Dyer in Hook. fil. Fl. Br. Ind. I, 298. *Mocanera grandiflora*, Blanco, Fl. Philipp. Ed. I, 451. *Anisoptera?* Turcz. in Bull. Soc. Nat. Mosc. 1858, I, 233.

Malacca: Maingay (Kew Distrib.) No. 198. Penang: Curtis 424. Perak: Scortechini 152 b. DISTRIB. Bangka, Teysmann. (?) Philip-pines.

The late Father Scortechini's field notes contain the following account of the flower: "The petals of this are red inside in the middle, but pale towards the margins; the stamens are numerous, 2-seriate, united in a ring by their enlarged bases, falling off together: staminodes many, short, adpressed to the ovary. Ovary pubescent, scaly towards the base. Fruiting-calyx reddish." The species comes near *D. Griffithii*: but is distinguished from it by the characters which I have noted under that species. Flowers of *D. Griffithii* are, however, wanting for comparison.

10. DIPTEROCARPUS KUNSTLERI, King, n. sp. A tree 80 to 120 feet high; young branches flattened, at first sparsely covered with minute scurfy deciduous pubescence, ultimately glabrous, but always with oblique tawny-tomentose annuli. Buds narrowly cylindric, hoary-canescenscent. Leaves elliptic or sub-rotund-elliptic, very shortly acaminate, the base rounded or sub-cuneate, the edges undulate or sub-crenate, both surfaces glabrous: main nerves 16 to 18 pairs, oblique, straight, prominent on the lower surface: length 7.5 to 11 in., breadth 4.5 to 7 in., petiole 1.5 to 2 in. Racemes 6 to 8 in. long, often bifid, 4- to 6-flowered, glabrous. Flowers 2.5 to 3 in. long, glaucous. Calyx-tube narrowly obconic, 5-winged, glaucous. Petals linear, obtuse, glaucous. Fruit sub-globular, an inch or more long, with 5 wings about .25 in. wide: accrescent calyx-lobes oblong, obtuse, slightly narrowed towards the base, glabrous, reticulate, 3-nerved, 6 or 7 in. long and about 1.25 in. broad.

Perak: King's Collector, Nos. 3638, 3798, 7508 and 7606.

Allied to *D. grandiflorus*; but with larger leaves, smaller fruit and different buds. Allied also to *D. Griffithii* but with smaller fruit and different buds. This species has leaves like *D. trinervis* Bl. and *D. retusus* Bl., but differs from these in having winged fruit: it also resembles *D. Dyeri*, Pierre, which, however, has longer leaves with hairy petioles and more narrowly winged fruit.

11. DIPTEROCARPUS GRIFFITHII, Miq. Ann. Mus. Lugd. Bat. I, 213. A tree 100 to 125 feet high: young branches stout, sub-compressed, minutely tawny-canescenscent; the leaf buds ovoid, densely covered with

yellowish-brown shining hair. *Leaves* coriaceous, broadly ovate, usually slightly narrowed to the rounded base, but sometimes the base truncate-ly sub-cordate, the apex acute or shortly acuminate, both surfaces glabrous, the upper shining; main nerves 12 to 14 pairs, spreading, straight, slightly prominent on the lower surface: length 5 to 11 in., breadth 3 to 5.5 in., petiole 2.25 to 3.5 in. *Racemes* 3- or 4-flowered. *Flowers* 1.5 in. long. *Calyx* ob-conic, sub-glabrous, 5-winged. *Fruit* oblong, 2.5 in. long, the wings extending from base to apex, stout, .5 in. or more broad: accrescent lobes of calyx oblong, obtuse, glabrous, reticulate, boldly 3-nerved, 5 to 7 in. long and about 1.75 in. broad. A. DC. in DC. Prod. XVI, Pt. 2, 611; Dyer in Hook. fl. Fl. Br. Ind. I, 299; Journ. Bot. 1874, 107. Kurz For. Flora Burm. I, 116. *D. grandiflorus* Griff. Notul. IV, 515 (not of Blanco).

S. Andaman: Kurz, King's Collector.

This closely resembles *D. grandiflorus*, Blanco, but the two may be readily distinguished by their young branches and leaf-buds. The young branches of this species are pale canescent and its leaf-buds broad and golden sericeous; while the branchlets of *D. grandiflorus* are quite glabrous and dark-coloured and the buds are narrow and pale canescent.

12. DIPTEROCARPUS INCANUS, Roxb. Hort. Beng. 42; Fl. Ind. II. 614. A tall tree: young shoots terete, stout, densely but minutely tawny-tomentose; the buds short, ovoid, thick, with longer tomentum than the branchlets. *Leaves* coriaceous, broadly ovate, acute or sub-acute, the base rounded or sub-cordate, the edges undulate; upper surface glabrous, the midrib alone slightly pubescent: under surface uniformly pale, shortly but softly stellate-pubescent, the midrib and nerves tomentose. main nerves 12 to 15 pairs, oblique, straight, prominent on the lower surface; length 5 to 8 in., breadth 2.5 to 4.75 in.; petiole .8 to 1.25 in., pubescent. *Flowers* about 1 in. long, usually in racemes but occasionally in short 7- or 8-flowered panicles. *Calyx-tube* ob-conic, 5-winged, minutely tomentose. *Petals* oblong, obtuse. *Fruit* sub-globose, about 1 in. in diam., 5-winged from base to apex; the wings thin, from .25 to .5 in. broad; the 2 accrescent lobes of the calyx narrowly oblong, obtuse, glabrous, much reticulate, 3-nerved in the lower half, when mature 5.5 in. long and nearly 1.5 in. broad; the 3 minor lobes sub-orbicular. Wight & Arn. Prod. 84; A. DC. Prod. XVI. 2, 611; Dyer in Hook. fl. Fl. Br. Ind. I, 298; Journ. Bot. 1874, p. 106.

S. Andaman: common. DISTRIB. Burmah, Kurz, Herb. No. 2109 (in part).

The plant here described under the name *D. incanus* closely re-

sembles *D. alatus*, Roxb.; but its flowers are shorter, the leaves are more broadly ovate, and have rounded or cordate, not cuneate, bases, while the pubescence of the lower surface is paler and more uniform and the young branchlets and leaf-buds are stouter. Moreover the accrescent lobes of the calyx are longer and nearly twice as broad: the 5 wings of the calyx-tube are also broader. Roxburgh's description of his species *D. incanus* is very brief; he left no drawing of it at Calcutta; and no authentic specimens of his own naming appear to exist. It is therefore impossible to decide with absolute certainty what Roxburgh's *D. incanus* is. At Kew Mr. Dyer accepts Kurz's Pegu specimen No. 2109 as belonging to it, and the specimens recently brought from the S. Andaman by my collectors agree with that number of Kurz's.

13. DIPTEROCARPUS ALATUS, Roxb. Hort. Beng. 42; Fl. Ind. II 614. A tree 80 to 125 feet high: young branches terete, rather stout, softly and minutely pubescent; the buds narrow, rufous-sericeous. *Leaves* coriaceous, ovate-elliptic, the apex acute, the base cuneate, the edges undulate: upper surface glabrous except the minutely tomentose nerves and midrib: lower sparsely and minutely stellate-pubescent, the 10 to 14 pairs of oblique rather straight prominent main nerves densely tomentose: length 5 to 8 in., breadth 2.75 to 4.5 in.; petiole 1 to 1.5 in., pubescent: stipules sericeous-pubescent. *Panicles* 6- or 7-flowered. *Flowers* about 1.5 in. long. *Calyx-tube* ob-conic, 5-winged, stellate-pubescent, as are the linear-oblong petals. *Fruit* globose, 1 in. in diam., puberulous, 5-winged from base to apex; the wings glabrous, thin and about .5 in broad; the 2 accrescent lobes of the calyx linear-oblong, obtuse, glabrous, much reticulate, 3-nerved in the lower half, 4.5 in. long and .7 or .8 in. broad: the 3 unenlarged lobes obtuse. Wall. Cat. 953; A. DC. Prod. XVI. 2, 611 *in part*: Dyer in Hook. fil. Fl. Br. Ind. I, 298; Journ. Bot, 1874, p. 106 (excl. syn. *D. costatus*, Gaertn.) Kurz For. Flora Burm. I. 116; Pierre Flore Forest. Coch-Chine, t. 212. *Oleoxyton balsamiferum* Wall. Cat. p. 157.

Burmah: Wallich, Brandis, Helfer No. 730, Kurz. Andamans?

Gaertner's figure and description of his *D. costatus* are confined to the fruit only. The former is that of a *Dipterocarpus* with the elongated calyx-lobes of *D. alatus*, Roxb., but with the 5 wings on the tube of the calyx very narrow, whereas those of Roxburgh's *D. alatus* are very broad. Dyer (F. B. I. i, 298) expresses his belief that Gaertner's figure is a bad representation of *D. alatus*, Roxb., and he reduces Gaertner's *D. costatus* to Roxburgh's *D. alatus*. M. De Candolle, on the other hand, retains *D. costatus*, Gaertn. as a good species and in this he is followed by Kurz; but Messrs. Dyer and De Candolle agree

that the *D. costatus* described by Roxburgh is a different plant from Gaertner's. For Mr. Dyer it is still a doubtful species; while M. De Candolle reduces it to *D. angustifolius* W. & A., which for Dyer is in its turn a doubtful species. A careful examination of the material now collected at Calcutta and at Kew leads me to believe that *D. costatus*, Gaertn., is a perfectly good species, and that the best character to distinguish it from Roxburgh's *D. alatus* is the narrowness of the wings of the calyx-tube. Specimens collected in Burmah by Kurz (No. 113 of his Herbm.) and by Brandis, have fruits exactly like that figured by Gaertner. Moreover I see no reason for thinking that the tree described by Roxburgh (Fl. Ind. II; 614) as *D. costatus*, Gaertn., is anything else than Gaertner's plant. Mr. Dyer (Journ. Bot. 1874, p. 153) expresses the opinion that *D. Lemeslei*, Vesque—a species collected on the island of Pulo Condor off the Cambodian coast—is reducible to *D. alatus*, Roxb.

It is very doubtful whether *D. alatus*, Roxb., occurs in the Andamans. I have seen no specimens of it from these islands, and I give it as an Andaman plant on the authority of the "Flora of British India."

Besides the preceding, there are various other species of *Dipterocarpus* in the Calcutta Herbarium from localities within the British Malayan region which, for want of sufficient materials, I am unable to describe. Chief amongst these are:—

- (1) Curtis No. 1560 from Penang, a species with winged calyx-tube.
- (2) A species from Perak, represented in Scortechini's collection (without number) by fruits resembling those of *D. Lowii* H., f., *D. intricatus*, Dyer, and *D. lamellatus*, Hook. fil.
- (3) A species from the Andamans with leaves resembling those of *D. Griffithii*, Miq., but with globular fruit which has neither angles nor wings on the calyx-tube. This possibly may be a form of *D. pilosus*, Roxb.
- (4) A Perak species (Herb. Scortechini mixed with No. 1478) represented by fruits something like those of *D. fagineus*, Vesque, but with the calyx-tube winged, not angled.
- (5) A Perak species represented by leaf-twigs and loose fruit of a species resembling both *D. fagineus*, Vesque, and *D. gracilis*, Bl., but differing from both.
- (6) A species from Perak (Wray No. 4031) having leaves like *D. Griffithii*, Miq., but with shorter petioles, and having also fruit rather like *D. Griffithii*, but the calyx-tube with narrower wings, and the minor calyx-lobes smaller.

2. ANISOPTERA, Korth.

Resinous trees. *Leaves* coriaceous, entire, feather-veined and finely reticulate; stipules small, fugacious or inconspicuous. *Flowers* in lax terminal panicles. *Calyx-tube* very short, adnate to the base of the ovary; the segments imbricate, then subvalvate. *Stamens* ∞ ; anthers ovoid with a long subulate connective, outer valves larger. *Ovary* 3- (rarely 4- 5-) celled; style fleshy, ovoid or oblong, with an attenuate 3-5-fid apex; ovules 2 in each cell. *Fruit* adnate to the calyx-tube, indehiscent, 1-seeded, crowned by the accrescent calyx-segments, of which 2 form linear-oblong lobes. *Cotyledons* fleshy, unequal; radicle superior. —DISTRIB. Malay Peninsula and Archipelago to New Guinea. Species about 6.

1. ANISOPTERA CURTISII, Dyer MSS. A tree 80 to 120 feet high: young branches slender, minutely scurfy-tomentose. *Leaves* oblong, tapering to both ends, the apex sub-acute or acute, the base narrowed but rounded; the upper surface glabrous, shining, the lower densely ochraceous-lepidote and sparsely stellate-pubescent; main nerves 18 to 20 pairs, spreading: length 2 to 3.5 in., breadth .75 to 1.25 in., petiole .5 to .75 in. Accrescent calyx-lobes 3.5 to 4.5 in. long, linear-spathulate, shining, 3-nerved: the transverse veins bold and numerous.

Penang: Curtis. Perak: King's Collectors.

Var. latifolia: leaves broadly elliptic, blunt, the bases rounded but narrowed.

Penang: Curtis, No. 1400.

The vernacular name of this in Penang is *Ringkong*.

3. VATICA, Linn.

Large or moderately sized resinous trees. *Leaves* coriaceous, entire, feather-veined and finely reticulate; stipules small, fugacious or inconspicuous. *Flowers* in axillary and terminal panicles, usually tomentose before expansion. *Calyx-tube* short, free, or adnate to the base of the ovary; segments somewhat acute, imbricate, then sub-valvate. *Stamens* 15; anthers oblong, external valves larger, connective apiculate. *Ovary* 3-celled; style short, subulate, or apex clavate or capitate; stigma entire or 3-toothed; ovules 2 in each cell. *Fruit* leathery, indehiscent, 1-seeded, surrounded by and sometimes partly adnate to the accrescent, membranous, nerved and reticulate calyx-lobes, two of which expand into narrow wings 2 or 3 in. long, the other three being much smaller. *Cotyledons* fleshy.

DISTRIB. Tropical Asia and chiefly Malaya; species about 10.

Synaptea is a genus established by Griffith (*Notulæ* IV., 516, Tab. 585 A, fig V.) for a tree collected at Mergui, and named by him *Synap-*

tea odorata. This plant has been named *Synaptea grandiflora* by Kurz, (Journ. A.S., Beng., 1870, 2, 65), and *Anisoptera odorata* Kurz, (For. Flor. Burm. I, 112), while Dyer has identified it with *Hopea grandiflora*, Wall, Cat. 958, and reduced it to *Vatica grandiflora* (F.B.I., i., 301).

The characters of the genus *Synaptea*, as given by its author, are practically those of *Vatica*, Linnæus (Mantissa II., p. 152-3, No. 1311), except that, whereas in the Linnæan description nothing is said about the fruit or its relation to the calyx, Griffith distinctly explains that he has given the name *Synaptea* because the ovary is adnate to the calyx. He does not say to what extent adnate, but, in fruiting specimens of his *Synaptea odorata*, the adhesion extends to the lower part only. In the "Mantissa" of Linnæus, only one species of *Vatica* is described, viz., *V. chinensis*; and of the specimen thus named in the Linnæan Herbarium, Sir J. G. Smith publishes a figure (Smith Ic., ined., t. 36.). This figure however does not show clearly whether the base of the ovary is, or is not, adherent to the calyx, and the fruit is not figured at all. A reference to Linnæus' specimen ought to settle what *V. chinensis* really is; but unfortunately it has not settled it. I have not myself examined the actual Linnæan specimen; but the opinions of botanists who have examined it vary as to its identity. The plant is generally admitted not to be of Chinese origin, for no Dipterocarp is known to inhabit China. Wight and Arnot are of opinion (Prod. 84) that *Vatica chinensis* is the same as *Vatica laccifera*, W. A. (*Shorea Talura*, Roxb.—*vide* Dyer). Alph. De Candolle (Prod. XVI., 2, p. 619) keeps up the species *V. chinensis*, while Dyer (Fl. Br. Ind., I, 302) reduces it to *Vatica Roxburghiana*, Blume (Mus. Bot. Lugd. Bat. II, 31. t. 7.), Blume's *Vatica Roxburghiana*, being, as the citations and figure given by that author show, the *Vateria Roxburghiana* of Wight's Illustrations, p. 87, and Icones t. 26. It cannot be demonstrated, therefore, either from Linnæus' description or specimen, or from Smith's figure of the latter, whether Linnæus intended his genus *Vatica* to include only plants with the ovary and fruit free from the calyx, or whether plants in which there is such partial adhesion might not also be admitted. If the latter were the case there would be no occasion to keep up the genus *Synaptea*. This is the view adopted by Messrs. Hooker and Bentham, who remark of *Synaptea*, "*ex descriptione auctoris verisimiliter ad Vaticam referenda est.*" This view is also adopted by Dyer, in "Hooker's Flora of British India," where he reduces *Synaptea odorata*, Griff., to the genus *Vatica*, Section *Eu-Vatica*. This view is also to a certain extent adopted by Burck who (Ann. Jard. Bot. Buitenzorg) makes *Synaptea* a section of *Vatica*, characterised by having the lobes of the fruiting

calyx unequally accrescent, two of them being much elongate, and *the fruit being partly inferior*; while the section *Eu-Vatica*, as proposed by Bentham and Hooker originally, and adopted by Burek, is characterised by having the same fruiting calyx as *Synaptea*; nothing being said about the adhesion between the calyx and the fruit. Pierre, on the other hand, keeps up *Synaptea* as a genus on account of the presence of albumen and the structure of the embryo (characters not easily worked in herbarium specimens of this family). In my own opinion it appears advisable to admit *Synaptea* as a section of *Vatica*, but to exclude *Isauxis*, *Retinodendron*, and *Pachynocarpus*, retaining these as distinct genera. *Vatica* would, according to this scheme, be divided into two sections:—

- I. *Eu-Vatica*:—Fruit free from the accrescent calyx, *i.e.*, fruit superior.
- II. *Synaptea*:—Fruit adnate in its lower part to the accrescent calyx, *i.e.*, fruit half inferior.
- Sect. I. EU-VATICA.—Fruit quite free from the calyx.
- Inflorescence and ripe fruit pale tomentose;
flowers 4 in. long 1. *V. perakensis*.
- Inflorescence and ripe fruit rusty-tomentose.
- Flowers 25 in. long; nerves of leaves 13
to 15 pairs; petioles 3 to 4 in. long ... 2. *V. Lowii*.
- Flowers 45 in. long; nerves of leaves 9
to 12 pairs; petioles 6 to 1.5 in. long... 3. *V. Maingayi*.
- Sect. II. SYNAPTEA.—Calyx-wings adherent to the
ripe fruit for nearly half its length.
- Leaves 9 to 10 in. long and with 18 to 20 pairs
of nerves 4. *V. nitida*.
- Leaves 2.5 to 7 in. long, with 6 to 13 pairs
of nerves.
- Larger lobes of calyx of fruit obovate and
very blunt.
- Leaves with 6 to 8 pairs of faint
nerves 5. *V. cinerea*.
- Leaves with 11 to 13 pairs of bold
nerves 6. *V. Curtisii*.
- Larger lobes of calyx narrowly oblong.
- Leaves oblong or elliptic-oblong,
with 9 to 11 pairs of nerves;
petals narrowly oblong ... 7. *V. faginea*.
- Leaves broadly elliptic, with 11 to 13
pairs of nerves; petals broadly
elliptic 8. *V. Dyeri*.

Leaves 2·5 to 3·5 in. long, with about 7 or 8 pairs of faint, main nerves, minutely reticulate. 9. *V. reticulata*.

1. *VATICA PERAKENSIS*, King, n. sp. A tree 60 to 80 feet high; young branches slender, deciduously scurfily stellate-pubescent, the bark rather pale. *Leaves* thinly coriaceous, oblong-lanceolate, rarely oblanceolate, more or less bluntly acuminate, sometimes caudate, the base cuneate; both surfaces glabrous, the midrib on the upper puberulous; main nerves 10 to 12 pairs, rather prominent beneath; length 2·5 to 4 in., breadth ·8 to 1·3 in., petiole ·4 to ·5 in. *Panicles* axillary and extra-axillary, crowded near the ends of the branches, 1 to 2 in. long, minutely pale tomentose, as are the ovate-lanceolate calyx-lobes. *Flowers* ·4 in. long. *Petals* narrowly oblong, obtuse, glabrous. *Stamens* slightly apiculate. *Ovary* minutely tomentose; stigma conical. *Ripe fruit* ·3 in. in diam., globose, the style persistent, minutely tomentose, quite free from the calyx; the two accrescent calyx-lobes oblong-ob-lanceolate, obtuse, obscurely 5-nerved, 2·5 in. long and ·5 in. broad; minor lobes unequal, lanceolate-acuminate, the largest about ·85 in long.

Perak: King's Collector, Wray; a common tree. Pangkore: Curtis.

The nearest ally of this is *Vatica Bantamensis*, Benth. and Hook.; but that has rather larger and more coriaceous leaves, which are perfectly glabrous; larger flowers with petals scaly externally and a more scurfy inflorescence; moreover the whole of the accrescent calyx-lobes of its fruit are more coriaceous and the minor lobes are blunter.

2. *VATICA LOWII*, King, n. sp. A tree 60 to 80 feet high: young branches, petioles, inflorescence and calyx densely rusty, scurfy-tomentose with stellate hair intermixed, the branches ultimately glabrous and with dark bark. *Leaves* coriaceous, oblong, sub-acute, the base rounded; both surfaces glabrous, the midrib puberulous on the upper; main nerves 13 to 15 pairs, spreading, slightly prominent beneath; length 2·5 to 3·5 in., breadth 1 to 1·5 in., petiole ·3 to ·4 in. *Panicles* axillary and terminal, much crowded towards the ends of the branches; ·75 to 1·5 in. long. *Flowers* ·25 in. long. *Calyx-lobes* lanceolate, acuminate, oblique. *Petals* narrowly oblong, obtuse, almost glabrous. *Stamens* short, unequal-sided, apiculate. *Ovary* depressed, tomentose, style capitate. *Ripe fruit* globular, ·25 in. in diam, deciduously rufous-scurfy; the style persistent, quite free from the calyx. Two large calyx-wings narrowly oblong, sub-acute, scarcely narrowed at the base, 5-nerved, 2·75 to 3 in. long, and ·6 in. broad; the three smaller lobes sub-equal, about ·5 or ·6 in. long, lanceolate, obtuse.

Perak: Scortechini, No. 2108; King's Collector, No. 7496.

This species is closely allied to *V. Maingayi*, Dyer; but has smaller flowers, and rather larger leaves with considerably longer petioles.

3. *VATICA MAINGAYI*, Dyer, in Hook. fil., Fl., Br., Ind. I, 302. A tall tree: young branches slender, ultimately glabrous, but at first rusty furfuraceous-tomentose, as are the inflorescence, calyx and ripe fruit. *Leaves* coriaceous, oblong or obovate-oblong, shortly acuminate, the base rounded, glabrous on both surfaces; main nerves 9 to 12 pairs, slender, curving, spreading; length 3 to 4·5 in., breadth 1 to 1·75 in., petiole ·6 to 1·5 in. *Panicles* short, few-flowered. *Flowers* ·45 in. long. *Calyx-segments* oblong-lanceolate. *Ovary* depressed, rufous-tomentose. *Ripe fruit* globose, ·25 in. in diam., the style persistent, rufous-tomentose; free from the calyx; the two large wings linear-oblong, sub-acute, not contracted at the base, 5-nerved (the lateral nerves faint) 2 in. long and ·35 to ·5 in. broad; the 3 smaller lobes ovate, sub-acuminate, ·75 in. long, all glabrous.

Malacca: Maingay (Kew Distrib.) No. 209.

Of this I have seen only Maingay's specimens, which are not good.

4. *VATICA NITENS*, King, n. sp. A tree 40 to 50 feet high: young branches and petioles densely covered with coarse deciduous scaly stellate tomentum, ultimately cinereous. *Leaves* coriaceous, narrowly oblong, acuminate, slightly narrowed to the rounded base; both surfaces, but especially the upper, shining, glabrous, the base on the lower sparsely scaly-tomentose when young, finely reticulate; main nerves 18 to 20 pairs, spreading, prominent on the lower surface: length 9 to 10 in., breadth 2 in.; petiole ·5 in., stout. *Ripe fruit* globular, crowned by the persistent style, reticulate, ·5 in. in diam., adnate for half its length to the calyx; the two large wings of the calyx oblong, slightly ob-lanceolate, obtuse, 3 in. long and ·8 to ·9 in. broad, the 3 shorter wings ovate-acuminate, ·8 in. long; all boldly 5-nerved and shining.

Penang: Curtis, No. 1404.

This fine species is known only by Mr. Curtis' imperfect specimens. It is very distinct, being at once recognisable amongst the Indian species of *Vatica* by the size of its leaves and calyx-wings.

5. *VATICA CINEREA*, King, n. sp. A tree about 40 feet high: young branches rufescent-puberulous at the very tips, otherwise glabrous and cinereous. *Leaves* thinly coriaceous, ovate-oblong to ovate-lanceolate, sub-acute, the base rounded or sub-cuneate; both surfaces glabrous, finely reticulate when dry; main nerves 6 to 8 pairs, spreading, faint; length 2·25 to 3·5 in., breadth ·75 to 1·5 in., petiole ·3 to ·5 in. *Panicles* mostly axillary, spreading, rusty scurfy-tomentose, 1·25 to 2 in. long. *Flowers* ·45 in. long. *Calyx-lobes* sub-equal, lanceolate, sub-acute, tomentose on both surfaces. *Petals* oblong-lanceolate, sub-acute, the half of the outer surface which is outside in æstivation pubescent, other-

wire glabrous. *Stamens* obtusely apiculate. *Ovary* depressed, minutely tomentose; stigma capitate. *Fruit* (not quite ripe) globular, umbonate, attached for half its length to the calyx. The two larger calyx-wings ob-lanceolate-oblong, obtuse or sub-acute, 5-nerved, flocculent-puberulous near the base when young, ultimately glabrous, 2 in. long and .5 in. wide; the 3 smaller wings lanceolate, obtuse, .5 in. long.

Langani: Curtis, Nos. 2797 and 2798. Kedah: Curtis, Nos. 2096 and 2514.

When dried, the leaves of this are of a dull gray colour—hence the specific name. Its fruit resembles that of the next species, but the leaves have fewer and less prominent nerves.

6. *VATICA CURTISII*, King, n. sp. A tree about 40 feet high: young branches, petioles, inflorescence and calyx brownish scurfy-pubescent, ultimately glabrous. *Leaves* ovate-oblong, sub-acute, the base rounded, both surfaces quite glabrous, reticulate; main nerves 11 to 13 pairs, oblique, rather prominent beneath; length 3 to 5 in., breadth 1.3 to 2.5 in., petiole .3 to .45 in. *Racemes* axillary, few-flowered, 1 to 1.25 in. long. *Flowers* .35 in. long. *Calyx-lobes* unequal, the 2 longer narrowly oblong, obtuse; the 3 shorter lanceolate-acuminate. *Petals* elliptic, slightly oblique, blunt, glabrous except the pubescent edge which is external in the bud. *Ripe fruit* globular, .3 in. in diam., adherent to the calyx for half its length, the larger calyx-lobes oblong-obovate, usually obtuse, rarely sub-acute, 5-nerved, 1.75 to 2 in. long, and .7 in. broad; the smaller wings about .4 in. long.

Penang: Curtis, No. 1579.

7. *VATICA FAGINEA*, Dyer in Hook. fl. Fl. Br. Ind., I., 301. A tree 80 to 100 feet high: young branches slender, minutely cinereous stellate-tomentose as is the inflorescence. *Leaves* coriaceous, oblong or elliptic-oblong, finely reticulate, glabrous; main nerves 9 to 11 pairs, spreading, curving, thin but prominent when dry; length 4 to 5 in., breadth 1.5 to 2 in. *Panicles* 2.5 in. long; flowers .5 in. long. *Calyx-tube* ribbed, minutely scurfy tomentose, the lobes unequal. *Petals* narrowly oblong, blunt, glabrous except the pubescent outside edge. *Ovary* hemispheric, minutely tomentose; stigma capitate, lobed. *Ripe fruit* globular, adherent for half its length to the calyx, about .25 in. in diam., the style persistent; the 2 larger calyx-wings narrowly oblong, or oblong-ob-lanceolate, obtuse, obscurely 5-nerved, 2 to 2.5 in. long, and .5 to .7 in. broad near the apex; the three smaller wings unequal, sub-spathulate, less than .5 in. long. *Hopea faginea*, Wall. Cat. 963 *Shorea pinangiana*, Wall., Cat. p. 157. *Synaptea faginea*, Pierre, For. Flore Coch.-Chine, t. 242.

Penang: Wallich. Perak: King's Collector, Nos. 3686 and 3765.

8. *VATICA DYERI*, King, n. sp. A tree 80 to 130 feet high: young branches, panicles, and calyx on both surfaces densely rufous-flocculent-tomentose, with stellate hairs intermixed, the branches ultimately glabrous and their bark pale. *Leaves* membranous, usually broadly elliptic, rarely elliptic-oblong, sub-acute or very shortly and bluntly acuminate, the base rounded, both surfaces quite glabrous, finely reticulate: main nerves 11 to 13 pairs, spreading, rather prominent beneath: length 3·5 to 7 in., breadth 1·6 to 3 in.; petiole ·35 to ·5 in., flocculent-tomentose. *Panicles* axillary or terminal, cymose, 1·5 to 3 in. long. *Flowers* ·4 in. long. *Calyx lobes* unequal, the two larger oblong and obtuse; the three smaller lanceolate, acuminate. *Petals* broadly elliptic, very obtuse, slightly narrowed to the truncate base, much larger than the calyx-lobes, glabrous, except one of the outside edges which is adpressed-pubescent. *Stamens* short, unequal-sided, bluntly apiculate. *Ovary* depressed-pubescent, the stigma capitate. *Ripe fruit* conical, the two large accrescent calyx-wings narrowly oblanceolate-oblong, blunt, 5-nerved, 1·25 in. long and ·25 in. broad; the three smaller wings one-fourth of the size of the larger, lanceolate, obscurely 5-nerved. *Synaptea Dyeri*, Pierre Fl. Forest. Coch-Chine, t. 241.

Perak: King's Collector, No. 7662. *DISTRIB.*, Cambodia, Lower Cochin-China, Pierre.

The Perak specimens are not in fruit: but in flowers and leaves they agree with Pierre's specimens from Cambodia and Cochin-China.

9. *VATICA RETICULATA*, King, n. sp. A tree 60 to 80 feet high: all parts except the inflorescence glabrous; young branches slender, dark-coloured. *Leaves* coriaceous, oblong to ovate-lanceolate, tapering from the middle to each end; the apex bluntly acuminate, the base very cuneate and slightly unequal-sided, the edges sub-undulate; both surfaces finely reticulate when dry, the lower paler; main nerves 8 or 9 pairs, little more prominent than the secondary; length 2·5 to 3·5 in., breadth 1 to 1·25 in., petiole ·4 in. *Panicles* axillary or terminal, puberulous, 2·5 to 3·5 in. long, lax, few-flowered. *Flowers* on long pedicels. *Calyx-lobes* unequal, lanceolate, more or less obtuse, densely pubescent on both surfaces. *Ovary* hemispherical, ridged, densely tomentose; style short, glabrous; stigma minute. *Young fruit* sub-globular; fruiting calyx with 2 accrescent linear-oblong wings, the other smaller; all attached to the lower part of the fruit.

Perak: King's Collector, No. 6969.

The only specimens which I have seen of this are without corolla, stamens, or ripe fruit. The species is, however, a very distinct one, and it is an unmistakable *Vatica*. I have therefore ventured to name it in spite of the imperfection of the material.

4. PENTACME, A. DC.

Glabrous or puberulous resinous trees. *Leaves* broad, entire, penninerved, with obtuse or cordate bases. *Flowers* large, paniced. *Calyx-tube* short, the lobes imbricate, 2 being quite external. *Stamens* 15, the filaments short, dilated; anthers much larger than the filaments, elongate, linear; the valves 4, sub-equal, each subulate at its apex, the connective also prolonged into a stiff deflexed arm as long as the appendages of the anther-valves. *Ovary* free; the style filiform, the stigma slightly lobed. *Fruit* enclosed within the imbricate calyx-lobes, of which two or more have elongated membranous reticulate many-nerved wings. Species 3,—Burmese, Siamese, and Malayan.

1. PENTACME MALAYANA, King, n. sp. A tree 40 to 50 feet high: young branches rather stout, dark-coloured, glabrous. *Leaves* subcoriaceous, rotund-ovate to broadly elliptic, the apex shortly and bluntly acuminate, the base rounded or slightly emarginate; both surfaces glabrous, pale when dry; main nerves 15 to 18 pairs, spreading, prominent on both surfaces; length 5 to 7 in., breadth 2·75 to 4·5 in., petiole ·75 to 1·1 in. *Panicles* axillary, lax, few-flowered, 2·5 to 5 in. long. *Flowers* ·75 in. long and about as much in diameter when open, pedicelled. *Calyx-lobes* more or less broadly ovate, acuminate, minutely tomentose outside. *Petals* three times as long as the calyx, elliptic, spreading, puberulous on one-half outside, and glabrous on the other, quite glabrous inside. *Stamens* 15, equal, erect, the filaments short and broad; the anthers elongate, narrow, with 5 apical awns, one of which is deflexed and rather shorter and thicker than the other four. *Ovary* ovoid, sub-glabrous, much shorter than the filiform style: stigma minute. *Ripe fruit* ovate, apiculate, 1 in. long, glabrous; calyx-wings all enlarged and reticulate except at the base; the three outer narrowly oblong, obtuse, and narrowed to the concave base, 9-nerved, 4 to 4·5 in. long, and ·65 to ·75 in. broad; the two inner lobes much narrower and fewer-nerved, about 2·5 in. long, or even shorter.

Langkani: Curtis, No. 2095.

The petals of this species are spreading, and the flower has quite an unusual *facies* for the order. It is at once distinguished by its curiously 5-awned anthers. Four of these awns are the produced apices of the anther cells, the fifth (the thicker and deflected one) is a prolongation from the connective.

5. SHOREA, Roxb.

Glabrous, mealy, or pubescent resinous trees. *Leaves* entire or sub-repand, pinnate-veined; stipules large, coriaceous and persistent, or minute and fugacious. *Flowers* in axillary or terminal, lax, cymose

panicles; bracts persistent, caducous, or 0. *Sepals* ovate or lanceolate, imbricate, 3 being external and 2 internal. *Stamens* 15 or 20, or 30; anthers ovate or oblong, rarely linear; connective subulate-cuspidate, rarely inappendiculate; valves obtuse, rarely cuspidate, equal, or the outer slightly larger. *Ovary* 3-celled, cells 2-ovuled; style subulate, stigma entire or 3-toothed. *Fruit* with leathery, rarely with woody, pericarp, 1-celled, 1-seeded, closely surrounded by the bases of the persistent, usually accrescent, sepals, the 3 outer, or more rarely, all, and sometimes none, of which are developed into 7- to 10-veined reticulate membranous linear-oblong wings. *Cotyledons* fleshy, unequal, usually enclosing the superior radicle. **DISTRIB**—Tropical Asia and chiefly the Malayan Archipelago: species about 60.

Sect. I. EU.-SHOREA. Fruit little more than .5 in. long, its pericarp leathery: three of the persistent sepals developed into membranous wings many times longer than the fruit.

Anthers without apical appendages.

Lower surface of adult leaves minutely stellate-tomentose, not scaberulous ... 1. *S. leprosula*.

Lower surface of adult leaves glabrescent, the axils of the nerves scaly ... 2. *S. scutulata*.

Lower surface of adult leaves quite glabrous, of young leaves glaucous ... 3. *S. Curtisii*.

Anthers mostly inappendiculate, a few with a minute apical appendage from the connective.

Stamens 30 4. *S. sericea*.

Anthers with very short apical appendages from the connective; flowers sessile.

Leaves 2.5 to 4 in. long, the lower surfaces minutely pubescent: flower .25 in. long; fruit ovoid-globose, its largest wings 2.5 in. long 5. *S. parvifolia*.

Leaves 3 to 4.5 in. long, glabrous beneath: flower .3 in. long; fruit turbinate, its largest wings 3.5 in. long ... 6. *S. acuminata*.

Leaves 4 to 6 in. long, glabrescent or glabrous beneath; fruit narrowly ovoid, its longest wings 3.5 to 4.5 in. long. ... 7. *S. macroptera*.

Apical appendage from the connective much longer than the anther.

Leaves glabrous on both surfaces, the lower not pale.

Stamens 10 (?) 8. *S. Maxwelliana*.

Stamens 20 9. *S. gratissima*.

Stamens 15

Flowers .2 to .25 in. long.

Main nerves of leaves 9 to 10 pairs,
faint; petals not saccate at base;
ovary ovoid-conical, tomentose,
style short 10. *S. Ridleyana*.

Main nerves 6 or 7 pairs; petals
saccate at base; ovary hemis-
pheric, style long and slender ... 8. *S. Maxwelliana*.

Flowers .4 in. long, main nerves 9 to
11 pairs; style 3 times as long
as the globose ovary 11. *S. pauciflora*.

Flowers .5 in. long, main nerves of
leaves 6 to 8 pairs; ovary elongate-
conic, style short, petals linear-
oblong 12. *S. Kunstleri*.

Flowers .65 in. long: nerves of leaves
12 to 16 pairs; ovary ovoid, style
long, filiform, petals ovate-lanceolate 13. *S. bracteolata*.

Leaves glaucous beneath 14. *S. glauca*.

Apical appendage of the connective with 3 to 5,
or many ciliae.

Stamens 30: ciliae radiating from the tip
of the apical process of all the anthers 15. *S. ciliata*.

Stamens 20: apical appendages of all the
anthers with numerous ciliae; petals
broad, spreading 16. *S. utilis*.

Stamens 15: anthers of outer row with
ciliate apical appendages 17. *S. costata*.

Anthers with a single apical appendage from
each cell, and a short one from the connec-
tive; sepals imbricate at their bases only ... 18. *S. stellata*.

Species imperfectly known.

Bracteoles large, persistent, scaberulous, stel-
late-pubescent 19. *S. Maranti*.

Stipules large, paired, persistent 20. *S. eximia*.

Sect. II. PACHYCHLAMYS, (Dyer). Fruit more than 1 in. long, its pericarp thick and woody, embraced in its lower half by a cup formed of the enlarged sepals, the bases of which are thickened woody and concave, the apices of the outer three produced into membranous wings as long as, or slightly longer than, the fruit.

Anthers of inner row inappendiculate, those

of the other two rows appendiculate ... 21. *S. Thiseltoni*.

1. *SHOREA LEPROSULA*, Miq. Fl. Ind. Bat. Suppl. I., 487. A tree 100 to 150 feet high: young branches rather slender, lenticellate, minutely and deciduously pale stellate-tomentose. *Leaves* coriaceous, elliptic to oblong, acute or sub-acute, the base rounded; upper surface glabrous, harsh from the prominent minute reticulations, the midrib and nerves sometimes puberulous; lower surface minutely fuscous-tomentose, with numerous densely stellate hairs on the midrib nerves and veins; main nerves 10 to 13 pairs, straight, oblique, prominent beneath; length 3 to 6 in., breadth 1·25 to 3·25 in., petiole ·35 to ·75 in. *Panicles* axillary and terminal, 1·5 to 4 in. long, rachis and branches stellate-tomentose, the short flower-bearing branchlets sericeous. *Flowers* in two rows, secund, 3 in. long, sessile. *Sepals* ovate, minutely velvety outside. *Petals* three times as long as the sepals, sericeous outside, oblong-spathulate. *Stamens* about 15; the filaments dilated, much longer than the short ovate inappendiculate anthers. *Ovary* ovoid, minutely tomentose, tapering upwards into the long slender style; stigma minute. *Ripe fruit* narrowly ovoid, apiculate, minutely tomentose, 6 in. long. *Calyx-wings* all enlarged and membranous, concave at the base so as to embrace the ripe fruit, but not adnate to it; the three outer narrowly oblong, sub-acute at the apex, narrowed at the base, 7-nerved, reticulate, 3 in. long and about ·7 in. broad; the two inner smaller, about 1 in. long, ovate, caudate-acuminate, not nerved. A. DC. Prod. XVI. 2, 631. Scheff. in Tijdschr. Ned. Ind. XXXI, 350: Hook. fil. Fl. Br. Ind., I., 305. Burck in Ann. Jard. Bot. Buitenzorg, VI, 215. *Shorea astrosticta*, Scortechini MSS.

Malacca: Maingay (Kew. Distrib.), No. 203. Perak, King's Collector, Nos. 7646, 7905, 8152; Scortechini, No. 2063. **DISTRIB.** Sumatra.

2. *SHOREA SCUTULATA*, King, n. sp. A large tree; young branches with dark lenticellate bark and minute white stellate pubescence. *Leaves* elliptic, shortly abruptly and bluntly acuminate; the base broad, rounded, almost truncate: upper surface glabrous, minutely reticulate; the lower, and especially the midrib, sparsely stellate-puberulous when young, glabrescent when old, the sides of the midrib, and especially the pits in the axils of the nerves, with numerous minute brownish pale-edged scales; length 3 to 3·5 in., breadth 1·5 to 1·75 in., petiole ·3 in. *Panicles* axillary and terminal, 3 to 4 in. long, the branches short, each bearing 2 or 3 bracteolate flowers; bracts broadly ovate, concave, blunt, hoary-puberulous, deciduous. *Flowers* 4 in. long, shortly pedicelled. *Sepals* broadly lanceolate, obtuse, tomentose outside, glabrous inside. *Petals* oblong, obtuse, the base expanded

at one side, glabrous inside and on one half outside, pubescent on the other. *Stamens* 15, in 3 rows; all the filaments broad, those of the outer two rows shorter than those of the inner: anthers short, broadly ovate, inappendiculate. *Ovary* conical, pale tomentose: style short, stigma small. *Fruit* (perhaps not mature) ovoid, apiculate, minutely pale tomentose, .6 in. long. *Sepals* all enlarged, membranous, reticulate, concave at the base; the three outer narrowly oblong, obtuse, very little narrowed to the base, 7-nerved, 2.75 in. long and .75 in. broad; the two inner .8 in. long, linear, about 1-nerved.

Penang: Curtis, No. 1396.

A species known only from Penang, and collected only by Mr. Curtis: remarkable for its almost racemose inflorescence, and curiously glandular leaves.

3. *SHOREA CURTISII*, Dyer MSS. in Herb. Kew. A tree 100 to 150 feet high; young branches slender, at first minutely stellate-puberulous, ultimately dark-coloured and glabrous. *Leaves* coriaceous, oblong-lanceolate, bluntly acuminate; the base sub-cuneate, or almost rounded; upper surface of young leaves minutely pubescent, of adults glabrescent or quite glabrous, the lower uniformly covered with very minute rufescent (young), or pale (adult) tomentum: main nerves 10 to 14 pairs, ascending, rather straight, prominent beneath: length 3 to 4 in., breadth 1.2 to 1.4 in., petiole .4 to .6 in. *Panicles* axillary or terminal, 2 to 3 in. long, the rachis slender, glabrous. *Flowers* about .3 in. long, in distichous secund rows of 4 or 5, on the short lateral branchlets, enveloped while in bud by broad deciduous puberulous bracts. *Sepals* ovate, tomentose outside, glabrous inside, slightly unequal. *Petals* twice as long as the calyx, linear-oblong, obtuse, stellate-pubescent outside, glabrous inside. *Stamens* 15, in three rows; the filaments elongate, broad (those of the outer row longest); anthers short, ovoid-globose, not apiculate. *Ovary* elongated ovoid, tomentose in the upper, glabrous in the lower half: style short, stigma small. *Ripe fruit* narrowly ovoid, apiculate, .75 in. long, pale tomentose; *calyx-wings* all enlarged and membranous, free from the fruit: the three outer linear-oblong, 8-nerved, 2.25 in. long, and about .5 in. broad; the two inner about 1 in. long, bluntly spatulate and with fewer nerves.

Penang: Curtis, Nos. 427, 1394 and 1395.

Perak: King's Collector, No. 8143.

The vernacular name of this in Penang is *Maranti Tai*.

4. *SHOREA SERICEA*, Dyer in Hook. fil. Fl. Br. Ind., I., 306. A tree 50 to 60 feet high; young branches rugulose, warted and scurfily

rufous-tomentose as are the inflorescence and petioles. *Leaves* coriaceous, oblong or elliptic-oblong (rarely slightly ob-ovate), very shortly acuminate or sub-acute, slightly narrowed to the rounded or sub-cuneate base; upper surface shining, sparsely stellate-tomentose, the depressed midrib and nerves puberulous; lower surface scaberulous, more densely stellate-pubescent, especially on the bold midrib and 20 to 22 pairs of stout spreading main nerves; length 3·5 to 6·5 in., breadth 1·5 to 2·75 in., petiole ·6 to ·8 in. *Panicles* axillary and terminal, 3 to 7 in. long, the ultimate branches bearing 4 or 5 distichous, secund, bracteate, sessile flowers; bracts broadly ovate, puberulous outside. *Sepals* ovate, the two inner smaller, all densely golden-sericeous outside, glabrous inside. *Petals* like the sepals and of about the same length, the inside and one-half of the outer glabrous, the other half adpressed-sericeous. *Stamens* about 40, in several rows; the filaments of the outer shorter, all longer than the anthers; anthers ovate, mostly inappendiculate, a few with a minute appendix. *Ovary* elongated, conic, sericeous; the style short, glabrous; stigma small. *Fruit* (immature) narrowly ovoid, ·5 in. long, embraced by, but not adnate to, the accrescent membranous calyx-wings: the outer 3 calyx-wings linear-oblong obtuse, narrowed to the base, 3·5 in. long and ·6 in. broad, 10-nerved; the 2 inner 2·5 in. long and much narrower and fewer-nerved, sparsely pubescent.

Malacca: Maingay (Kew. Distrib.) No. 202. Penang: Curtis, No. 431. Perak: King's Collector, No. 3511.

This resembles *S. lacunosa* Scheff., but differs in not having persistent stipules. Its vernacular name in Penang is *Seraya*.

5. *SHOREA PARVIFOLIA*, Dyer in Hook. fil. Fl. Br. Ind., I., 305. A tree 100 to 150 feet high; young branches slender, pale tomentose at first, ultimately glabrous, dark-coloured and lenticellate. *Leaves* coriaceous, ovate to ovate-lanceolate, caudate-acuminate, the base sub-cuneate or almost rounded; upper surface glabrous (when young the midrib tomentose or pubescent); under surface sparsely scaly-pubescent when young, when adult minutely pubescent, the transverse veins thick; main nerves 9 to 12 pairs, oblique, rather straight, prominent beneath: length 2·5 to 4 in., breadth 1 to 1·8 in.; petiole ·35 to ·45 in., tomentose when young. *Panicles* axillary and terminal, crowded near the ends of the branches, 2 to 4 in. long, rather lax, spreading, many-flowered, minutely tomentose, the branches distichous. *Flowers* ·25 in. long, secund, distichous, deciduously bracteate. *Sepals* slightly unequal, ovate, acute, tomentose outside, glabrous inside. *Petals* twice as long as the sepals, obliquely elliptic, obtuse, glabrous, except on one-half outside which is silky. *Stamens* 15, or fewer: the filaments flatten-

ed, about 4 times as long as the broad short anthers; apiculus of connective very slender, about as long as the anther, deflexed. *Ovary* elongate, puberulous; style rather short; stigma small. *Ripe fruit* ovoid-globose, .4 in. long, thinly adpressed pale tomentose. *Sepals* all enlarged and membranous, concave at the base so as to embrace the ripe fruit, but not adnate to it: the three outer narrowly oblong, obtuse at the apex, slightly narrowed to the base; 7-nerved, 2.5 in. long; the two inner from one-half to one-third shorter, narrower and fewer nerved. *Shorea disticha*, Scortechini MSS. in Herb. Calcutta.

Malacca: (Kew Distrib.) No. 206. Penang: Curtis, No. 201. Perak: Scortechini, No. 1965. Wray, No. 1282.

6. *SHOREA ACUMINATA*, Dyer in Hook. fil. Fl. Br. Ind., I., 305. A tree 100 to 150 feet high; young branches minutely greyish tomentose, ultimately dark-coloured and glabrescent. *Leaves* coriaceous, ovate to lanceolate, acuminate, the base often unequal-sided, rounded or sometimes emarginate; upper surface glabrous except the puberulous midrib; the flower glabrous, with a few scattered stellate hairs: main nerves 7 to 9 pairs, spreading, slightly prominent beneath: length 3 to 4.5 in., breadth 1.75 to 2.5 in.; petiole .3 to .4 in., tomentose. *Panicles* axillary and terminal, crowded near the extremities of the branches, 2 to 3 in. long, minutely stellate-pubescent, many-flowered. *Flowers* .3 in. long, distichous, secund, about 5 on each lateral branch, bracteolate. *Sepals* ovate, unequal, tomentose outside, glabrous inside. *Petals* twice as long as the calyx, spreading, broadly ovate, puberulous outside, glabrous inside. *Stamens* 15, in three rows, the inner row shorter: filaments broad, much larger than the short, ovate, minutely appendiculate anthers. *Ovary* ovoid, tapering, pubescent: style short, stigma small. *Ripe fruit* turbinate, with 3 slightly vertical grooves, apiculate, puberulous, .5 in. in diam., attached by its base to the calyx: *sepals* all enlarged, concave at the base so as completely to cover the fruit, membranous and reticulate; the 3 outer narrowly oblong obtuse, contracted towards the base, 10- or 11-nerved, 3.5 in. long, and 7 in. broad; the two inner 1 to 1.5 in. long, under .25 in. broad, 3- to 4-nerved.

Malacca: Maingay (Kew Distrib.) No. 205 (?). Griffith, No. 1762. Perak: King's Collector, No. 8009.

7. *SHOREA MACROPTERA*, Dyer in Hook. fil. Fl. Br. Ind. I, 308. A tree 60 to 80 feet high: young branches with dark-brown bark, minutely lenticellate and puberulous. *Leaves* coriaceous, oblong (usually narrowly), shortly acuminate, the base sub-cuneate or rounded: upper surface glabrous, shining, the midrib and nerves puberulous: lower

surface glabrescent or glabrous, chocolate-coloured when dry: main nerves 10 to 12 pairs, curved, spreading, prominent on the lower surface; length 4 to 6 in., breadth 1·35 to 1·75 in.; petiole ·4 to ·5 in., rugose. *Panicles* axillary or terminal, 4 to 7 in. long, lax, branching, few-flowered, puberulous, sparsely scaly. *Flowers* about ·5 in. long, sessile, solitary, not secund. *Sepals* distinct almost to the base, slightly unequal, broadly-ovate, acute, more or less yellowish-tomentose outside, glabrous inside. *Petals* narrowly oblong, slightly oblique at the base, the apex blunt, glabrous except one-half of the outer surface which is sericeous. *Stamens* 15, in two rows; filaments broad except at the apex, those of the outer two rows by much the shorter: anthers short, ovate, the connective minutely awned. *Ovary* elongated-ovoid, sericeous in its upper half; style short, stigma small. *Ripe fruit* ·6 to ·75 in. long, narrowly ovoid, pale puberulous, apiculate: *sepals* all enlarged and reticulate, slightly concave at the base and embracing, but not adnate to, the fruit; the three outer narrowly oblong, obtuse, tapering slightly to the auricled base, 7-nerved, 3·5 to 4·5 in. long, and ·8 to 1 in. broad; the two inner variable, but shorter, narrower and fewer nerved. *Shorea auriculata*, Scortechini MSS. in Herb., Calcutta.

Malacca: Maingay. Singapore: Ridley. Penang: Curtis, No. 1392. Perak: very common, King's Collector, Scortechini.

A species from Borneo which closely resembles this appears to me to differ specifically. Its leaves are longer with sparser nerves, and its calyx-wings are longer.

8. *SHOREA MAXWELLIANA*, King, n. sp. A tree 60 to 80 feet high: young branches dark-coloured, almost glabrous. *Leaves* coriaceous, ovate-lanceolate, acuminate (caudate-acuminate when young), the base unequal-sided, cuneate; both surfaces quite glabrous, the upper shining, the lower chocolate-coloured when dry: main nerves 6 or 7 pairs, curved, spreading, thin and inconspicuous: length 3 to 4 in., breadth 1·3 to 1·5 in., petiole ·4 in. *Panicles* axillary and terminal, 2·5 to 3 in. long, stellate-puberulous, their lateral branches very short and few-flowered. *Flowers* shortly pedicelled. *Sepals* unequal, oblong, blunt, with enlarged concave bases, more or less pubescent, but glabrous in the concavity of the base inside. *Petals* oblong, concave and saccate at the base, tomentose outside, glabrous inside. *Stamens* 10 (?), the filaments short, broad; the anthers elongate, erect, pointed, the connective ending in an awn as long as the anther. *Ovary* hemispheric; the style long, slender; stigma minute. *Fruit* (not mature) globular, minutely tomentose, closely invested by, but not adnate to, the concave bases of the sepals: *sepals* all enlarged, membranous, narrowly oblong, obtuse;

the three outer 7-nerved, 1·5 in. long and 4 in. broad; the two inner similar in shape, but fewer-nerved and only 5 in. long.

Perak: King's Collector, Nos. 3601 and 3744.

The only flowers of this species which I have seen are in an early stage of bud, and from them I am unable to make out the characters of the petals properly. The stamens appear to be only 10 in number: but of this I cannot now be quite certain.

9. *SHOREA GRATISSIMA*, Dyer in Hook. fil. Fl. Br. Ind. I, 307. A tree: younger branches slender, glabrescent, dark-coloured. *Leaves* coriaceous, elliptic, acuminate, the base broad and rounded, the margins sub-undulate, both surfaces glabrous: main nerves 12 to 14 pairs, faint; length 2·5 to 4 in., breadth 1·25 to 1·5 in., petiole 6 to 75 in. *Panicles* axillary and terminal, lax, few-flowered, 3 to 6 in. long, sub-puberulous. *Flowers* secund, pedicelled, 25 in. long. *Sepals* lanceolate, sub-acute; minutely tomentose outside, glabrous inside in the lower, adpressed-pubescent in the upper, half. *Petals* twice as long as the calyx and much broader, elliptic, obtuse, glabrescent. *Stamens* about 20; the filaments short, unequal, dilated. *Anthers* elongated-ovate, truncate, each with a terminal awn from the connective twice as long as itself. *Ovary* ovoid, sub-glabrous; stigma small. *Ripe fruit* unknown. *Hopea gratissima*, Wall. Cat. 960.

Singapore: Wallich.

This is known only by Wallich's specimens. He referred it to *Hopea*, of which genus it certainly has the *facies*: the aestivation of the sepals is moreover that of *Hopea*, and so is the apiculus of the connective of the stamens. The petals in shape, however, resemble those of *Shorea*. I retain it in *Shorea* in deference to the opinion of Mr. Dyer.

10. *SHOREA RIDLEYANA*, King, n. sp. A tree 60 or 80 feet high: young branches slender, dark brown, lenticellate, nearly glabrous. *Leaves* ovate-lanceolate, shortly acuminate, the base rounded: both surfaces glabrous, the upper shining: main nerves 9 or 10 pairs, curved, spreading, thin but slightly prominent beneath: length 2·5 to 4 in., breadth 1·1 to 2 in.; petiole 4 to 5 in., rugulose. *Panicles* axillary and terminal, 1·5 to 2 in. long, densely stellate-puberulous. *Flowers* 2 in. long, pedicellate. *Sepals* sub-equal, oblong, obtuse, tomentose outside, glabrous inside. *Petals* oblong, slightly oblique, obtuse, glabrous inside, puberulous outside on one half, glabrous on the other. *Stamens* 15, sub-equal, the filaments dilated in the lower half: anthers shorter than the filaments, ovate, the connective produced into an awn longer than the anther. *Ovary* ovoid-conical, minutely tomentose. *Style* short; stigma minute. *Fruit* (immature) ovoid, apiculate, minutely

pale tomentose: *sepals* all enlarged, membranous, reticulate and concave at the base; the three outer linear-oblong, obtuse, slightly narrowed to the concave base, 5-nerved, 2·25 in. long and ¼ in. broad; the two inner of the same shape, but only 1-nerved, narrower and only 1·5 in. long.

Perak: King's Collector, Nos. 3571 and 3617.

This a good deal resembles *S. Maxwelliana*, King; but its leaves have more nerves, its slightly oblique petals are not saccate at the base, its ovary is ovoid-conical, and minutely tomentose with a short style; whereas in *S. Maxwelliana* the petals are saccate at the base, and the ovary is hemispheric with a long style.

11. *SHOREA PAUCIFLORA*, King, n. sp. A tree 50 to 90 feet high: young branches slender, their bark brown puberulous and lepidote. *Leaves* thinly coriaceous, from oblong to elliptic, shortly acuminate; the base abruptly cuneate, slightly unequal-sided, or (in the elliptic forms) almost rounded: main nerves 9 to 11 pairs, oblique, straight, prominent beneath: length 4 to 5 in., breadth 1·8 to 2·5 in., petiole ¼ to ⅞ in. *Panicles* few, axillary or terminal, few-flowered, 1·75 to 4 in. long, rather coarsely pubescent. *Flowers* ¼ in. long, secund, shortly pedicellate, each subtended by an ovate, solitary, puberulous, deciduous bract. *Sepals* broadly ovate, tomentose outside, glabrous inside. *Petals* broadly elliptic, obtuse, concave at the base, veined, inside glabrous, the outside half glabrous and half adpressed-sericeous. *Stamens* 15, in 3 rows: the outer row smaller and with filiform filaments, the inner rows with filaments longer and expanded in the lower half; the anthers of all shortly ovate, the connective produced into an awn twice as long as the stamen. *Ovary* hemispheric, tomentose; style nearly 3 times as long, puberulous; stigma small. *Ripe fruit* unknown.

Penang: Curtis, No. 1537.

A species known only by Mr. Curtis' specimens which have no fruit.

12. *SHOREA KUNSTLERI*, King, n. sp. A tree 60 to 100 feet high: young branches slender, rusty-puberulous, their bark brown. *Leaves* coriaceous, elliptic, abruptly and shortly acuminate, the base rounded or slightly cuneate, both surfaces glabrous, the lower with a few stiff white hairs on the midrib and nerves; main nerves 6 to 8 pairs, curved, ascending, prominent on the lower surface; length 4 to 5 in., breadth 2 to 2·4 in., petiole ⅝ in. *Panicles* axillary and terminal, 4 to 6 in. long, lax, few-flowered, scaly-puberulous. *Flowers* ⅝ in. long, sub-sessile, 4 or 5 together on the short branches of the panicles, secund, bracteate: the bracts broadly ovate, puberulous. *Sepals* sub-equal,

broadly ovate, acute, tomentose outside; the edges ciliate, glabrous inside. *Petals* linear-oblong, obtuse; the bases obliquely expanded, sericeous externally, glabrous internally. *Stamens* 15, sub-equal, the filaments as long as the anthers, flattened; anthers ovate, short, the connective terminated by a curved awn much longer than the stamen. *Ovary* elongate-conic, puberulous; style short. *Ripe fruit* hemispheric, tapering into a cone and crowned by the style, adpressed pale tomentose. *Sepals* membranous, reticulate: the three larger narrowly oblong, obtuse, tapering to the concave non-reticulate base, 9-nerved, 3·5 in. long and 7 in. broad: the two inner 2 in. long, linear, 3-nerved.

Perak: King's Collector, Nos. 3474 and 3705.

This species is allied to *S. bracteolata*, Dyer, but its leaves have fewer nerves, smaller flowers, narrower petals, and a short style.

13. *SHOREA BRACTEOLATA*, Dyer in Hook. fil. Fl. Br. Ind. I, 305. A tree 50 to 150 feet high; young branches minutely furfuraceous-puberulous, speedily glabrescent, their bark dark-coloured. *Leaves* coriaceous, elliptic-oblong, shortly acuminate (often sub-obtuse when old), narrowed slightly to the rounded or emarginate base; upper surface quite glabrous; the lower yellowish furfuraceous-puberulous to glabrous; main nerves 12 to 16 pairs, spreading, prominent beneath: length 4 to 6 in., breadth 1·6 to 2·5 in., petiole 45 to 6 in. *Panicles* axillary, few-flowered, 2·5 to 6 in. long, glabrous. *Flowers* 65 in. long, shortly pedicellate, each subtended by 2 elliptic, obtuse, 3-nerved, puberulous, deciduous bracts 35 in. long. *Sepals* lanceolate, obtuse, minutely tomentose outside, the two inner smaller. *Petals* ovate-lanceolate, obtuse; the bases expanded, glabrous. *Stamens* 15, in two rows, the filaments less than half as long as the ovate obtuse anthers; appendix of connective subulate, twice as long as the anther, decurved when old. *Ovary* ovoid, attenuated upwards, sub-glabrous; the style long, filiform; stigma small. *Ripe fruit* ovoid, apiculate, 6 in. long, embraced by, but (except at the very base) free from the calyx; *sepals* accrescent, membranous, reticulate and concave at the base: the three outer narrowly oblong, blunt, slightly narrowed above the concave base, 10-nerved, 3·5 in. long, and 6 in. broad; the two smaller about 2 in. long, and 2 in. broad, about 3-nerved. *Shorea foveolata*, Scortechini MSS. in Herb. Calcutta.

Malacca: Maingay (Kew Distrib.) No. 204. Penang: Curtis, Nos. 322 and 1405. Perak: King's Collector, Nos. 7583, 7591, 7717; Scortechini, No. 1939. **DISTRIB.**—Sumatra. Forbes, No. 3050.

14. *SHOREA GLAUCA*, King, n. sp. A tree 80 to 100 feet high; young branches slender, dark-coloured, puberulous. *Leaves* coriaceous, ovate-lanceolate, acuminate; the base broad, rounded; upper surface

glabrous, the lower glaucous (except the midrib and nerves) especially when young; main nerves 7 to 9 pairs, ascending, rather straight: length 3·5 to 4·5 in., breadth 1·4 to 1·8 in.; petiole ·45 to ·6 in., rugulose, glaucous. *Panicles* axillary, few-flowered, shorter than the leaves, hoary, the *flowers* on short pedicels. *Sepals* slightly unequal, oblong, obtuse, tomentose on both surfaces. *Ovary* conical, tomentose; the style very short, glabrous; stigma small, 3-lobed. *Fruit* (immature) ovoid-globose, apiculate, minutely tomentose; accrescent sepals membranous, free from the fruit; obscurely 7- to 12-nerved, strongly reticulate, blunt, slightly narrowed to the concave base, at first puberulous but ultimately glabrous; the longer 2·25 in. long, and ·6 to ·75 in. broad, the others smaller.

Penang: Curtis, No. 372. Malacca: Maingay (Kew Distrib.), 212.

In this species the two inner fruiting wings of the calyx are nearly as large as the three outer; the leaves are very white underneath when young, but much less conspicuously so when adult. It is known, only by Curtis' and Maingay's specimens, none of which have complete flowers. Maingay's specimens from Malacca have in fact no flowers; but there is no mistaking their leaves as being exactly like those of Mr. Curtis' from Penang. The vernacular name of this is *Dammar laut dhan lesor*.

15. *SHOREA CILIATA*, King, n. sp. A medium-sized tree; young branches slender, dark-coloured, deciduously hoary-puberulous. *Leaves* coriaceous, lanceolate or oblong-lanceolate, acuminate, the base cuneate; both surfaces glabrous, minutely reticulate, the lower whitish when young, pale brown when dry; main nerves 8 or 9 pairs, ascending, curved, shining on the lower surface: length 3 to 3·5 in., breadth ·8 to 1·5 in., petiole ·75 to ·9 in. *Panicles* 2 to 2·5 in. long, axillary and terminal, little-branched, few-flowered, hoary. *Flowers* ·5 in. long, secund. *Sepals* ovoid-deltoid, obtuse, outside tomentose, inside glabrous. *Petals* three times as long as the sepals, narrowly oblong, obtuse, slightly expanded at the base, adpressed-sericeous outside, glabrescent inside. *Stamens* 30, in fascicles of 3, unequal, the shorter with undilated filaments, the longer with filaments dilated in the lower half; all with the connective produced into an apical process crowned by 3 to 5 spreading ciliæ. *Ovary* ovoid-conic, sericeous, with a short glabrous style. *Fruit* (immature) ovoid, apiculate, pale-tomentose, ·5 in. long; accrescent sepals membranous, reticulate: the three outer narrowly oblong, reticulate, 7-nerved: the two inner 2 in. long, and ·3 in. broad, narrowed to above the concave base: the two inner 1 in. long, linear-lanceolate, few-nerved.

Penang: Curtis, No. 1578.

Known only by Curtis' specimens, and readily recognisable by its beautifully ciliate-crested anthers.

16. *SHOREA UTILIS*, King, n. sp. A large tree; all parts except the inflorescence glabrous: young branches slender, dark-coloured. *Leaves* coriaceous, ovate-lanceolate, caudate-acuminate, or shortly and abruptly acuminate, the base slightly cuneate; main nerves about 7 pairs, oblique, not prominent on either surface; length 2·5 to 3 in., breadth ·9 to 1·2 in., petiole ·4 in. *Panicles* axillary, stellate-puberulous, about as long as the leaves; their lateral branches distant, very short, minutely tomentose, 3- or 4-flowered. *Flowers* sub-sessile, globular in bud, under ·2 in. long. *Sepals* ovate-orbicular, blunt, the outer 3 very tomentose outside, the inner 2 less so; all glabrous inside. *Petals* broadly oblong, blunt, more or less sericeous in both surfaces. *Stamens* 20; filaments slightly dilated, about as long as the ovate anthers; apical process of connective about as long as the anther, ciliate. *Ovary* sericeous, elongated-conic, gradually tapering into the short glabrous style; stigma minute. *Ripe fruit* ovoid, apiculate, pale, adpressed-sericeous, ·4 in. long, closely invested by, but free from, the concave bases of the accrescent sepals. *Sepals of fruiting calyx* all enlarged, membranous, reticulate, deciduously puberulous; the 3 outer oblong, very obtuse, 5-nerved, 1·25 in. long, and ·4 in. broad; the inner 3 half as long, or less, and much narrower.

Penang: Curtis, No. 423.

This species, which Mr. Curtis describes as yielding the most durable timber in Penang, was at one time quite common there, but it is now almost extinct. Its vernacular name is *Dammar laut*.

17. *SHOREA COSTATA*, King, n. sp. A tree; young branches dark-coloured, lepidote-puberulous. *Leaves* thinly coriaceous, oblong, sub-acute, slightly narrowed to the rounded or sub-cuneate base; both surfaces glabrous, the transverse veins distinct, especially on the lower: main nerves 11 to 13 pairs, oblique, rather straight, slightly prominent beneath; length 3 to 4·25 in., breadth 1·2 to 1·5 in., petiole ·8 to 1 in. *Panicles* axillary and terminal, 1·5 to 2·5 in. long, scaly-puberulous, the lateral branches very short and few-flowered. *Flowers* small. *Sepals* broadly ovate, yellowish-tomentose outside, glabrous inside. *Stamens* 15; all with dilated filaments longer than the ovate anthers, those of the inner row with the apical process of the connective short and glabrous, those of the outer rows with longer ciliate apical connectives. *Ovary* ovoid-conical, densely yellowish-tomentose; style very short. *Ripe fruit* ovoid, apiculate, sparsely puberulous, ·75 in. long; sepals all enlarged, concave and dilated at the base, membranous and reticulate; the three outer narrowly oblong, obtuse, much

narrowed to the base, 7-nerved, 2·75 in. long, and ·45 in. broad; the two inner of the same shape, but few-nerved, only 1·5 in. long, and ·25 in. broad.

Penang: Curtis, No. 199.

A species known only by Mr. Curtis' solitary specimen. The connectives of the inner anthers are ciliate, somewhat in the fashion of *S. ciliata*, King; but the leaves of that species are very different.

18. *SHOREA STELLATA*, Dyer in Hook. fil. Fl. Br. Ind. I, 304. A tree 100 to 150 feet high; young branches slender, at first stellate-puberulous, but speedily glabrous, with bark dark-coloured and sparsely lenticellate. *Leaves* thinly coriaceous, ovate-lanceolate, the base rounded: upper surfaces glabrous, the lower very minutely lepidote on the reticulations; main nerves 8 to 11 pairs, rather straight, oblique, prominent on the lower surface; length 4 to 5·5 in., breadth 1·75 to 2·25 in., petiole ·7 to ·9 in. *Panicles* axillary or terminal, crowded at the extremities of the branches, many-flowered, 4 to 6 in. long; minutely stellate-pubescent. *Flowers* ·25 in. in diam. *Calyx* minutely greyish-tomentose, the segments ovate-oblong, sub-acute, valvate, erect. *Petals* broadly ovate, obtuse, pubescent outside, spreading. *Stamens* 15, the filaments short, broad; the anthers linear-elongate, shortly bi-mucronate, the connective also shortly mucronate. *Ovary* ovate-globular, grooved, very tomentose; the style short; the stigma ovoid, small. *Ripe fruit* ovoid, apiculate, tomentose, ·5 in. long; sepals all enlarged, sub-equal, membranous, linear-oblong, sub-acute, much narrowed at the base, quite free from the fruit, 5-ribbed, reticulate, 4·5 in. long, and about ·6 in. broad. *Parashorea stellata*, Kurz, Journ. As. Soc., Bengal, for 1870, pt. 2, p. 66. For. Flora Burm., I, 117; Pierre Flore Forest. Coch-Chine, t. 224.

Perak: King's Collector, No. 7505. DISTRIB. Burmah.

None of the Perak specimens are in fruit; but in leaves and flowers they agree absolutely with Kurz's Burmese specimens. The calyx in all is quite valvate, and it was on this character chiefly that Kurz based the genus *Parashorea*.

19. *SHOREA MARANTI*, Burck in Ann. Jard. Bot. Buitenzorg, VI. 217. A small tree: young branches dark-coloured, stellate-puberulous. *Leaves* thinly coriaceous, more or less broadly elliptic or elliptic-oblong, shortly abruptly and bluntly acuminate; the base broad, rounded, or almost truncate; upper surface glabrous, the midrib and nerves minutely tomentose or pubescent when young; lower surface more or less sparsely minutely stellate-puberulous, the sides of the midrib, especially at the axils of the main nerves, glandular and densely covered with masses of brown pale-edged scales: main nerves 12 to 16 pairs, oblique,

slightly curved, thin but prominent beneath when dry, as are the transverse veins; length 3·5 to 6·5 in., breadth 1·5 to 2·25 in.; petiole ·35 in., densely stellate-pubescent, scurfy. *Stipules* deciduous, ovate-lanceolate, nerved, stellate-puberulous. *Panicles* axillary and terminal, few-flowered, tawny-tomentose, (shorter than the leaves [?]); the bracts in pairs, unequal, elliptic-oblong, blunt, nerved, pubescent on both surfaces. “Segments of calyx (fide Burck) unequal, the three outer larger, imbricate. *Petals* minutely tomentose inside. *Stamens* 15, in two rows.” *Hopea? Maranti*, Miq. Fl. Ind. Bat. Suppl., 489; A. DC. Prod. XVI, 2, p. 635.

Perak: King’s Collector, No. 880. Malacca: Derry, No. 952. **DISTRIB.** Sumatra, Bangka.

The Perak specimens are not in flower; and I have seen none from elsewhere that are. The above imperfect description of the flower has therefore been copied from Burck (Ann. Jard. Bot. Buitenzorg, VI. 217). The Perak specimens perfectly agree, as to leaves, with an authentic specimen of Miquel’s from Sumatra, in the Calcutta Herbarium. Miquel never saw either flower or fruit. In fact, of the twenty new species of *Dipterocarpeæ* described by this author in the supplement to his Flora of the Netherlands India, the flowers are described in only two, and in these but partially!

20. *SHOREA EXIMIA*, Scheff. in Nat. Tijdschr. Ned. Ind. XXXI, 349. A shrub or small tree; young branches petioles and under-surfaces of leaves stellate-setulose. *Leaves* coriaceous, elliptic-oblong, or ob-lanceolate-oblong, acuminate, narrowed to the rounded or sub-cuneate base: upper surface glabrous except the tomentose midrib, shining, the nerves depressed: under surface scabrid, pale brown, the reticulations midrib and 17 to 21 pairs of spreading nerves prominent: length 6·5 to 11 in., breadth 2·25 to 3·25 in., petiole ·25 to ·35 in. *Stipules* in pairs, persistent, ovate, acuminate, longer than the petioles, reticulate, laxly pubescent and warted. “*Wings of fruiting-calyx* linear-lanceolate, obtuse: the three larger narrow at the base, 3·2 to 3·6 in. long, ·5 in. broad, sparsely pubescent, 9-nerved; the two shorter and narrower 1·6 in. long. *Fruit* elongated-ovoid, acuminate, minutely whitish-tomentose.” Burck in Ann. Jard. Bot. Buitenzorg VI, 218. *Vatica? eximia*, Miq. Fl. Ind. Bat. Suppl. 486; A. DC. Prod. XVI, 2, 623. *Vatica sub-lacumosa? Miq.* Fl. Ind. Bat. Suppl. 486. *Shorea sub-lacumosa*, Scheff. in Nat. Tijdschr. Ned. Ind. XXXI, 350: A. DC. Prod. XVI, 2623.

Malacca: Griffith, No. 5018. Penang: King. Perak King’s Collector, 10998. **DISTRIB.** Sumatra, Bangka.

This plant is very imperfectly known. I have copied the descrip-

tion of the fruit from Dr. Burek (l. c.). Miquel, who first described the plant as a probable *Vatica*, had seen nothing but a leaf-twig. Specimens brought from Perak by the Calcutta collectors bear, instead of flowers, curious cones, 1·5 in. long, of distichous imbricate bracts, concerning which Griffith, in his field note on his specimen No. 5018, wrote,—“irregular growth caused by an insect; each of the scales of these cones bears on its dorsum at its base a number of eggs.” Griffith’s No. 5019 appears to belong to a closely allied, but distinct, species; as also does the indeterminate plant issued by Wallich as No. 6635 of his catalogue, under the designation, “*Dilleniaceae* [?] *nervosa*.”

21. *SHOREA THISELTONI*, King, n. sp. A tree 60 to 80 feet high: young branches rather stout, the bark dark-coloured and lenticellate, but covered at first by a pale-grey, deciduous pellicle. *Leaves* coriaceous, elliptic-oblong to elliptic, rarely oblong, sometimes slightly obovate, obtuse, slightly narrowed to the rounded base; both surfaces glabrous, the lower when very young sparsely lepidote, puberulous especially on the midrib and nerves, brown when dry: main nerves 8 or 9 pairs, ascending, slightly curved, bold and prominent on the under surface like the midrib; length 5 to 7 in., breadth 2·5 to 3·25 in.; petiole ·6 to ·8 in., stout. *Panicles* axillary and terminal, 2 to 3 in. long, velvety, few-flowered, apparently ebracteolate. *Flowers* sessile, ·6 or ·7 in. long. *Sepals* ovate, sub-acute, unequal; the 3 outer tomentose outside, glabrous inside; the 2 inner smaller, nearly glabrous, the edges ciliate. *Petals* much longer than the sepals, linear-oblong, obtuse, expanded at the base, glabrous, except one-half of the outer surface which is adpressed-pubescent. *Stamens* 15, in 3 rows, the filaments of all dilated, unequal: the anthers shortly ovate, those of the inner and longer row inappendiculate, those of the other two rows with a short apical appendage from the connective. *Ovary* narrowly conical, tomentose, tapering into the short glabrous style; stigma minute. *Type fruit* narrowly ovoid, apiculate, minutely pale-tomentose, substrate, 1·2 in. long, and ·6 in. in diam., the pericarp thick and woody. *Persistent sepals* with much thickened concave woody bases, forming a cup embracing the lower half of the fruit, the apices of the outer three prolonged into membranous linear-oblong obtuse wings exceeding the fruit and sometimes 1·5 in. long; one of the inner sepals shortly winged, the other often broad, obtuse and not winged.

Perak: common. King’s Collector.

In this plant the fruit is much larger than in any of the other species of *Shorea* here described, and its pericarp is hard and thick. The bases of the sepals are greatly thickened and concave, and they form a cup which embraces closely, but does not adhere to, the lower

half of the fruit, the apices of some of them being winged as above described. In these respects the species resembles certain other Malayan species of *Shorea*, e. g., *S. Martiniana* Scheff, *S. scaberrima*, and *S. stenoptera*, Burck. Judging from the leaf-specimens on which Miquel founded his *Hopea Singkawang*, that plant must be a close ally of this. A species (flower only) collected by H. O. Forbes in Sumatra (Herb. No. 2952) must also be closely allied to this. It differs however by its conspicuously bracteolate inflorescence. Beccari's Nos. 2681 and 3507, which form the types of Heim's species *S. brachyptera*, are also allied to this.

6. HOPEA, Roxb.

Glabrous or hoary-tomentose resinous trees. *Leaves* quite entire, firm, feather-veined; stipules small, deciduous or inconspicuous. *Flowers* sessile or shortly pedicelled, ebracteate, in lax panicles of unilateral racemes. *Sepals* inserted on the receptacle, two being quite external and three for the most part internal, obtuse, imbricate. *Petals* falcate, their apices inflected in bud. *Stamens* 15, or rarely 10, slightly connate; the connective subulate-cuspidate, the anthers ovate, their valves obtuse, equal. *Ovary* 3-celled, the cells 2-ovuled: style shortly cylindrical or subulate. *Fruit* 1-seeded, closely surrounded by the bases of the accrescent sepals, the 2 external of which are developed into linear wings, the three internal not longer than the ripe fruit. *Embryo* as in *Shorea*.—DISTRIB. of *Shorea*; species about 35.

Sect. I. EU-HOPEA, Main nerves of leaves bold and prominent.

Nerves of leaves 16 to 18 pairs; accrescent sepals 4 to 4.5 in. long, 10-nerved ... 1 *H. nervosa*.

Nerves of leaves 10 to 13 pairs; accrescent sepals 1.75 to 2.5 in. long, obscurely 5-nerved 2. *H. Curtisii*.

Sect. II. DRYOBALANOIDEA, Miq. Main nerves not distinct.

Petals sericeous: the filaments longer than the anthers; ripe fruit .3 in. long, the accrescent sepals 7-nerved, 1.75 to 2 in. long, and .2 to .25 in. broad; leaf-petioles .25 to .4 in. long, minutely tomentose... 3. *H. micrantha*.

Petals densely sericeous; the filaments shorter than the anthers; ripe fruit .2 in. long; accrescent sepals obscurely 5- to 7-nerved, 1.25 to 1.5 in. long, and .25 in. broad; leaf-petioles .35 to .6 in. long, slender, puberulous, finally glabrous ... 4. *H. intermedia*.

1. *HOPEA NERVOSA*, King, n. sp. A tree 50 to 70 feet high: young branches dark-coloured, glabrous. *Leaves* coriaceous, oblong to elliptic-oblong, shortly acuminate, the base rounded or very slightly cuneate; both surfaces glabrous; main nerves 16 to 18 pairs, spreading, bold and prominent on the lower; length 3·5 to 5 in., breadth 1·5 to 2·25 in.; petiole ·5 to ·75 in., transversely wrinkled when dry. *Flowers* unknown. *Ripe fruit* ovoid-rotund, apiculate, glabrous, ·5 in. long; the two outer sepals much enlarged, oblong-lanceolate, obtuse, slightly narrowed to the concave thickened smooth base, 10-nerved, 4 to 4·5 in. long, and ·6 to ·75 in. wide; the three inner sepals not quite so long as the fruit, broadly ovate, obtuse, thickened, smooth, closely embracing but not adherent to the fruit.

Perak: King's Collector, No. 3690.

This is a very distinct species, belonging to the group of *Hopea* with the nerves of the leaves bold. It is so distinct that, contrary to my general practice, I venture to name it without having seen the flower.

2. *HOPEA CURTISII*, King, n. sp. A tree 50 to 60 feet high: young branches slender, dark-coloured, lenticellate, almost glabrous. *Leaves* coriaceous, broadly ovate to ovate-oblong, shortly acuminate or acute, the base slightly unequal-sided, rounded, rarely sub-cuneate; both surfaces glabrous, the upper slightly puberulous on the midrib near the base, the lower with several hairy glands at the base, the midrib sparsely and minutely stellate-puberulous; main nerves 10 to 13 pairs, curving, ascending, prominent beneath; length 3·5 to 4·5 in., breadth 1·75 to 2·5 in.; petiole ·4 in., puberulous when young. *Panicles* axillary and terminal, lax, few-flowered. *Flowers* about ·2 in. long, pedicelled. *Sepals* broadly ovate, blunt, concave, tomentose outside, glabrous inside; the inner two rather smaller and more glabrous than the others. *Petals* oblong, oblique, falcate, obtuse, partially tomentose outside, glabrous inside. *Stamens* 10, the filaments short, dilated; anthers ovate, short, the connective with an apical awn longer than the anther. *Ovary* broadly ovate, puberulous at the truncate apex, otherwise glabrous: style short. *Ripe fruit* ovoid, apiculate, pale striate, ·3 in. long; outer two sepals accrescent, narrowly-oblong, reticulate, membranous, obscurely 5-nerved, obtuse, slightly narrowed to the concave smooth base, 1·75 to 2·5 in. long and from ·35 to ·6 in. broad; the three inner non-acrescent sepals about as long as the fruit.

Penang: Curtis No. 1562. Perak: King's Collector, 8161.

3. *HOPEA MICRANTHA*, Hook. fil. in Trans. Linn. Soc., xxiii, 160. A tree 60 to 80 feet high: young branches slender with dark-coloured, lenticellate bark and minute brownish pubescence. *Leaves* coriaceous,

ovate-lanceolate or oblong-lanceolate, bluntly caudate-acuminate; the base slightly cuneate or sometimes broad, rounded and slightly unequal; both surfaces glabrous except the pubescent midrib: main nerves numerous, not much more prominent than the secondary, and both indistinct; length 2 to 4 in., breadth .8 to 1.75 in.; petiole .25 to .4 in. minutely tomentose. *Panicles* axillary and terminal, numerous, short, spreading, 1 to 1.5 in. long, puberulous or glabrous. *Flowers* .15 to .25 in. long, shortly pedicellate. *Sepals* sub-equal, ovate-rotund, sub-acute or obtuse, puberulous and resinous outside, glabrous inside. *Petals* twice as long as the sepals, broadly oblong-obtuse, silky outside except on one side, glabrous inside. *Stamens* about 12, the filaments dilated in the lower half, longer than the ovate anthers; the connective produced into a single apical awn longer than the stamen. *Ovary* elongated, often constricted in the middle, glabrous; style very short, stigma minute. *Ripe fruit* ovoid, apiculate, .3 in. long, striate, closely embraced by the 3 inner sepals which about equal it in length; the outer two sepals accrescent, oblanceolate, obtuse, tapering to the concave base, reticulate, 7-nerved, 1.75 to 2 in. long, and .2 to .25 in. broad. A. DC. Prod. XVI. 2, p. 634. Dyer in Hook. fil. Fl. Br. Ind. I, 310. Burck in Ann. Bot. Jard. Buitenzorg, VI, 238.

Malacca; Maingay (Kew Distrib.) No. 210. Penang: Curtis, Nos. 167, 266, 1397. Perak: King's Collector, Nos. 3525, 8170. DISTRIB. Borneo: Bangka, Sumatra.

Mr. Curtis notes on the Penang specimens of this, that the bark of the tree is smooth and of a grey colour, whereas the bark of its close ally *H. intermedia* is fissured like that of *Shorea parviflora*. The species of *Hopea* with numerous indistinct nerves, (Sect. *Dryobalanoides*) are not easy to distinguish from each other in the Herbarium. *H. Mengarawan*, Miq., a species published two years earlier than this (*i. e.*, in 1860), comes very near this, and the two may possibly prove to be identical, in which case Miquel's name must be adopted. *Hopea cernua*, Teysm. and Binn. was described by its authors from a plant originally obtained from Sumatra, but cultivated in the Buitenzorg Garden. It differs from *H. Mengarawan* and from *H. micrantha* in having larger leaves with more prominent nerves. Its authors were doubtful as to its being really distinct from *H. Mengarawan*, and I think these doubts were well founded. Under the species named *H. Dryobalanoides* by Miquel (*l. c.*) there are, Dr. Burck asserts, two plants. One of these collected at Soengie-pagoe in Sumatra, is, he says, simply *H. Mengarawan*, Miq., and it is the fruit of this which Miquel describes under his *H. Dryobalanoides*. The other specimen from Priaman in Sumatra is different, and it is to it that Dr. Burek (Ann. Bot. Jard. Buitenzorg VI., 241) desires to

restrict the name *H. Dryobalanoides*, Miq. There is in the Calcutta Herbarium an authentic specimen of the very gathering of the Soengie-pagoe plant on which Miquel worked, and I should refer it to *H. micrantha* Hook. fil.

Petalandra micrantha, Harssk. has been reduced by the authors of the *Genera Plantarum* (Vol. I. p. 193) to *Hopea*. It is however a different plant from this, and belongs to Miquel's section *Eu-hopea*, which is characterised by the nerves being prominent. By Dr. Burck, *Petalandra* is reduced to *Doona*.

4. HOPEA INTERMEDIA, King n. sp. A tree 60 to 80 feet high : young branches rather dark-coloured, minutely lenticellate, puberulous. *Leaves* coriaceous, ovate-lanceolate, caudate-acuminate, the base cuneate, both surfaces glabrous; main nerves numerous, faint; length 2·5 to 3 in., breadth 1 to 1·35 in.; petiole ·35 to ·6 in. slender, puberulous but finally glabrous. *Panicles* as in *H. Mengarawan*, the flowers pedicellate. *Sepals* sub-equal; the two outer ovate, acuminate; the three inner broader and more obtuse, all resinous outside, glabrous and smooth inside. *Petals* twice as long as the sepals, narrowly oblong, obtuse, falcate, densely sericeous externally, glabrous within. *Stamens* 12; the filaments dilated, shorter than the anthers; the anthers short, crowned by a straight awn from the connective longer than the stamen. *Ovary* hour-glass shaped; style short, stigma small. *Ripe fruit* ovoid, apiculate, ·2 in. long, pale, striate; the two outer sepals accrescent, narrowly oblong-obtuse, narrowed to the base, reticulate, obscurely 5- to 7-nerved, 1·25 to 1·5 in. long and ·25 in. broad; the inner three sepals not accrescent, not longer than the fruit, and closely embracing it.

Penang: Curtis, No. 425 and 1398. Perak: King's Collector, No. 3709.

This species is no doubt near to *H. micrantha*, Hook. fil., but, according to Mr. Curtis, it is distinguishable from that, while growing, by its bark, this tree having a fissured bark like that of *Shorea parvifolia*, Dyer, while the bark of *H. micrantha* is smooth and grey. The petals of this are also more sericeous than those of *H. micrantha*, the filaments are shorter than the anthers (not longer, as in *H. micrantha*), the leaves are more glabrous, the petioles longer and more slender and more glabrous, and the fruit and accrescent sepals are smaller than in *H. micrantha*. I have therefore ventured, after much hesitation, to name this as a species, and from its relationship to *H. micrantha* and *H. Mengarawan*, I have called it *H. intermedia*. Its vernacular name in Penang is *Jankang*. It has been suggested that this plant should be referred to *H. Dryobalanoides*, Miq.—a course which I would have adopted with great pleasure had it been clear what *H. Dryobalanoides* really is.

But, as I have stated in a note under *H. micrantha*, *H. Dryobalanoides* appears to be a composite species; moreover, its author nowhere describes its flowers. For these reasons I think it ought to be suppressed as a species.

7. RETINODENDRON, Korthals.

Resinous trees, with the leaves, inflorescence, and flowers of *Vatica*. *Ripe fruit* globular, crowned by the persistent style, 1-celled, 1-seeded, the pericarp coriaceous, indehiscent. *Calyx* of ripe fruit slightly accrescent, the pieces oblong, nearly equal, and quite free from, and usually shorter than, the fruit (longer in *R. Kunstleri*). *Isauxis* (sub-genus of *Vateria*) W and A. DISTRIB. Malaya and British India. Species about 10.

Isauxis was established by Wight and Arnot as a sub-genus of *Vateria*, Linn. to receive the three species *Vateria lanceæfolia*, Roxb., *V. Roxburghiana*, Wight and *V. Ceylonica*, Wight (*Stemoporus Wightii*, Thw.) and its characters were, "Segments of the calyx ovate, acute, enlarging in fruit; petals falcate and about three times the length of the calyx: *stamens* 15 with oblong anther cells; *style* short; *stigma* clavate, 3-6 toothed: *panicles* axillary, shorter than the leaves." The other section of *Vateria* suggested by Wight was *Eu-Vateria* (the *Vateria* of Linnæus and of which *V. indica*, L. is the type) and of this the characters are, "*Calyx-segments* obtuse, scarcely enlarging in fruit: *petals* oval, scarcely longer than the calyx: *stamens* 40 or 50 with linear anther-cells: *style* elongated: *stigma* acute; *panicle* large and terminal. Korthals, evidently overlooking Wight's Illustrations, published (Verh. Nat. Gesch. Ned. Ind. p. 56) his genus *Retinodendron* to cover one of the very plants (*viz.*, *Vateria lanceæfolia*, Roxb.) for which Wight and Arnot founded the sub-genus *Isauxis*; and to this *Retinodendron* Korthals added his own Malayan species *R. Rassak* and *R. pauciflorum*. Although *Isauxis* may have the priority as a sub-genus (Wight's Illustrations were published in 1840, and Korthals' book, just quoted, bears the date 1839-1842), *Retinodendron* takes precedence as a *genus*. The flowers of *Retinodendron* are exactly those of all the species of *Vatica* (except the anomalous *V. scaphula*, Roxb.) inasmuch as the segments of the calyx are slightly imbricate when the bud is very young, becoming valvate as the bud advances in age; the petals are much longer than broad, their apices are not inflexed in aestivation, and they are not spreading when expanded. The fruit itself is also practically that of *Vatica*; but the fruiting-calyx is different, for its lobes are invariably free from the beginning, they are pretty nearly equal to each other, but (although slightly accrescent) they are in most cases *shorter* than the fruit. As regards its calyx, *Retinodendron* is closely allied to *Vateria*, but it differs from *Vateria* in its flowers; for in *Vateria* the stamens are numerous (40 to 50), the petals are scarcely longer than the segments of the calyx and are spreading; moreover the inflorescence is longer in *Vateria* than in *Retinodendron*, and it is terminal. In short, *Retinodendron* has the flowers of *Vatica* and the fruit of *Vateria*. Dr. Burck forms *Retinodendron* and *Isauxis* into sections of the genus *Vatica*, giving however characters to the section *Isauxis* which form no part of Wight's original characters of it as a sub-section of *Vateria*. In Dr. Burck's section *Isauxis*, "the calyx-lobes are all accrescent, sub-equal to the fruit, or much longer."

Fruiting-calyx shorter than the fruit.

Leaves 3·5 to 6 in. long : fruit ¼ in. in diam. 1. *R. pallidum*.

Leaves 7 to 10 in. long : fruit ⅙ in. in diam. 2. *R. Scortechinii*.

Fruiting-calyx longer than the fruit ... 3. *R. Kunstleri*.

1. *RETINODENDRON PALLIDIUM*, King. A small tree (fide Dyer) : young branches slender, deciduously puberulous, their bark pale. *Leaves* coriaceous, oblong-lanceolate to narrowly elliptic, acuminate ; the edges entire, recurved when dry ; the base acute : both surfaces glabrous, the upper shining ; main nerves 9 to 10 pairs, curving, oblique ; length 3·5 to 6 in., breadth 1·2 to 1·8 in., petiole ¼ to ⅕ in. *Panicles* axillary, rarely extra-axillary, puberulous, 1 to 3 in. long. *Flowers* ¼ in. long ; *Calyx-segments* ovate-lanceolate, scurfy-pubescent. *Petals* oblong, lanceolate, sub-acute, stellate-pubescent externally. *Anthers* broadly ovate, with a short blunt apiculus. *Ovary* puberulous ; stigma capitate, lobed. *Fruit* globular, about ¼ in. in diam., glabrous, shining, very minutely and sparsely lepidote, partially covered in the lower half by the slightly unequal, spreading or sub-reflexed, narrowly-oblong, membranous, 3-nerved, reticulate calyx-lobes. *Vatica pallida*, Dyer in Hook. fil. Fl. Br. Ind. I, 302.

Penang : Maingay, on Government Hill, at an elevation of about 800 feet ; Curtis, No. 117 ; King, Kunstler.

This is known only from Penang. It is evidently a rare tree. Its fruit somewhat resembles (except in size) that of *V. lanceæfolia*, Blume.

2. *RETINODENDRON SCORTECHINII*, King, n. sp. A tall tree : young branches rather stout, densely furfuraceous-pubescent. *Leaves* coriaceous, oblong, tapering to the sub-acute apex ; the base slightly narrowed, rounded : both surfaces glabrous : main nerves 14 to 18 pairs, spreading, curving, prominent on the lower, depressed on the upper, surface when dry, the transverse venation bold : length 7 to 10 in., breadth 2·6 to 3·2 in., petiole ⅙ to ⅚ in. *Panicles* crowded towards the apices of the branches, mostly axillary, 2 to 2·5 in. long, the rachises brownish flocculent stellate-tomentose, as is the calyx externally. *Flowers* ⅙ in. long. *Calyx-lobes* ovate. *Petals* thick, oblong, blunt, puberulous externally, glabrous within. *Stamens* elliptic, apiculate. *Ovary* minutely tomentose ; stigma clavate. *Ripe fruit* sub-globular, sub-rugose, vertically grooved, minutely rufous-scurfy, about ⅙ in. in diam., laxly embraced in the lower half by the broadly ovate, membranous, many-nerved, reticulate, sub-equal calyx-lobes.

Perak : Scortechini, Nos. 1940 and 1942.

The calyx-lobes are nearly equal in size, quite free from the fruit, much shorter, and they embrace only its lower half. This species is allied to *Retinodendron Rassak*, Korth. (Nat. Gesch. Ned. Ind. 56, t. 8.)

but has broader leaves and much more condensed panicles than that species.

3. *RETINODENDRON KUNSTLERI*, King n. sp. A tree, 20 to 50 feet high, sometimes a shrub: young branches slender, deciduously stellate-puberulous. *Leaves* thinly coriaceous, elliptic-oblong to oblong-lanceolate, sometimes slightly obovate, sub-acute or shortly and bluntly acuminate; the base cuneate, rarely rounded: upper surface glabrous, the midrib and nerves pubescent; the lower quite glabrous; main nerves 7 to 9 pairs, ascending, slightly prominent beneath: length 2·25 to 4·5 in., breadth 1·25 to 1·75 in., petiole ·25 to ·4 in. *Racemes* axillary, 1 to 1·5 in. long, sparsely scaly. *Flowers* ·4 in. long. *Calyx-lobes* ovate-lanceolate, puberulous. *Petals* oblong-elliptic, oblique, obtuse, puberulous outside. *Anthers* slightly and sharply apiculate. *Ovary* puberulous, stigma capitate. *Ripe fruit* globular, with a long curved apical beak, glabrous, about ·25 in. in diam. *Calyx-lobes* all accrescent, sub-equal, oblong, tapering slightly to the sub-obtuse apex, the base slightly auricled, thickly membranous, glabrous, 3-nerved, the longest about 1·3 in. long, and ·35 in. broad, loosely surrounding, and longer than, the fruit.

Perak; Scortechini, Wray, King's Collector; very common at low elevations.

In this species all the five calyx-lobes are accrescent and of nearly equal size. They are quite free from the ripe fruit, round which they form a loose semi-inflated investiture. Its nearest ally is *Vatica bancana*, Scheffer, (*Retinodendron bancanum*).

8. ISOPTERA, Scheffer.

A tall resinous tree. *Leaves* coriaceous, entire, feather-veined. *Flowers* in axillary or terminal panicles. *Calyx-tube* very short, the segments ovate-rotund, imbricate. *Stamens* 30 to 35, the anthers ovate, the cells divergent at the base, acute, the valves equal, the connective produced into an apical bristle-like appendage. *Ovary* 3-celled, the loculi bi-ovulate; the style short, terete, 3-angled at the apex. *Fruit* indehiscent, 1-seeded, the pericarp coriaceous. *Fruiting-calyx* an open cup not embracing the fruit; its lobes all slightly enlarged, spreading (not winged); the outer 3 rotund, broader than the 2 narrower inner lobes.

One species—Malayan.

1. *ISOPTERA BORNEENSIS*, Scheff. MSS. ex Burck in Ann. Bot. Jard. Buitenzorg VI, 222. A large tree: young branches slender, dark-coloured, sparsely lenticellate, glabrescent. *Leaves* coriaceous, oblong, sub-acute, slightly narrowed to the rounded base: upper surface glabrous except the puberulous midrib; the lower pale, glabrous; main

nerves 8 or 9 pairs, oblique, slightly curving, prominent beneath; length 4 to 5 in., breadth 1.75 to 2 in., petiole .5 in. *Panicles* 4 to 6 in. long, stellate-pubescent; bracteoles caducous. *Flowers* shortly stalked. *Ocalyx-segments* minutely tomentose. *Petals* .5 in. long, pale tomentose. *Stamens* 30 to 36, in 3 series, the filaments dilated at the base: anthers with equal valves. *Ovary* sericeous, style glabrous. *Ripe fruit* subglobose, acuminate, pale tomentose, about .25 in. in diam.; fruiting-calyx forming a cup with a concave short tube embracing the fruit, the segments spreading, re-curved, the 3 outer .65 in. in length and breadth, the 2 inner smaller. Heim, "Recherches sur les Dipterocarpacees," p. 51.

Pahang: Ridley, No. 2626. **DISTRIB.** Bangka, Borneo.

Leaf-specimens of what appear to be this tree were collected by Mr. Wray (Herb. No. 3426) in Upper Perak.

9. *BALANOCARPUS*, Beddome.

Glabrous or puberulous, rarely scabrid, resinous trees, with inconspicuous fugaceous stipules. *Leaves* entire, coriaceous or membranous, penni-nerved. *Flowers* secund, sessile or shortly pedicelled. *Sepals* distinct or united at the base, imbricated, two quite external to the others; in fruit sub-equal, only slightly enlarged, woody, thickened, and forming a 5-lobed cup round the base of (but rarely enveloping) the fruit, not adnate to it and never expanding into wings. *Petals* elliptic, obliquely acuminate, the apices slightly inflexed in bud or not inflexed at all. *Stamens* 15, attached to the bases of the petals, in 3 rows; or 10 in 2 rows, sub-equal, the filaments much dilated at the base, the connective prolonged into a straight apical awn longer than the ovate anther. *Torus* flat. *Ovary* 3-celled, cells 2-ovuled, ovules collateral. *Style* short. *Stigma* minute, entire. *Fruit* oblong or sub-globose, apiculate; the pericarp ligneous or sub-ligneous. *Seed* solitary, erect; cotyledons fleshy, plano-concave, the larger 2- or 3-lobed, or entire; the radicle prominent. Southern Peninsular India, Malaya. Probably 12 species.

Leaves glabrous, smooth.

Leaves ovate-lanceolate or ovate, caudate-acuminate.

Stamens 15

Fruit entirely enveloped in the slightly enlarged calyx

1. *B. Curtisii*.

Only the lower part of the fruit enveloped by the calyx

2. *B. penangianus*.

Stamens 10

3. *B. anomalus*.

Leaves narrowly oblong, gradually narrowed to the acute apex.

Fruit 1·75 to 2·25 in. long : stamens 10 ... 4. *B. maximus.*

Fruit 1·5 in. long ; leaves 4 to 6 in. long, with 9 or 10 pairs of bold parallel nerves 5. *B. Heimii.*

Fruit ·6 in. long : leaves 2·25 to 2·75 in. long, with 7 or 8 pairs of slightly prominent nerves 6. *B. Wrayi.*

Leaves stellate-pubescent, scabrid 7. *B. Hemsleyanus.*

1. *BALANOCARPUS CURTISII*, King. A tree 20 to 30 feet high : young branches slender, the bark dark-coloured, puberulous. *Leaves* membranous, ovate-lanceolate, bluntly caudate-acuminate, the base slightly cuneate : both surfaces glabrous, dull ; main nerves 8 to 10 pairs, spreading, faint and scarcely more prominent than the secondary nerves ; length 2 to 2·5 in., breadth ·75 to 1 in., petiole ·1 to ·15 in. *Panicles* axillary and terminal, shorter than the leaves, glabrescent, lax, each with a few 3- to 5-flowered spreading branches. *Flowers* secund, shortly pedicelled, ·15 in. long. *Sepals* distinct, sub-equal, thick, rotund-ovate, very obtuse, puberulous outside, glabrous inside, the edges slightly ciliate. *Petals* elliptic, obliquely shortly and bluntly acuminate, glabrescent inside, partly puberulous and partly glabrous outside. *Stamens* 15, in 3 rows, sub-equal ; the filaments shorter than the anthers, dilated : anthers broadly elliptic, truncate, the connective produced into an apical awn longer than the stamen. *Ovary* cylindric, truncate, glabrous, the style short and stigma minute. *Fruit* smooth, globular, apiculate, crowned by the sub-sessile discoid stigma, enveloped by, but not adherent to, the slightly thickened sepals, ·25 to ·3 in. in diam. (calyx included).

Penang : Curtis, No. 1406. Perak : King's Collector, Nos. 3171, 3294, 6543 ; Wray, No. 2860.

2. *BALANOCARPUS PENANGIANUS*, King, n. sp. A tree 40 to 50 feet high : young branches slender, dark-coloured, lenticellate, slightly puberulous at the very tips. *Leaves* coriaceous, ovate-lanceolate or ovate-acuminate, often caudate-acuminate, the base slightly cuneate or almost rounded, the edges slightly undulate, both surfaces glabrous : main nerves 7 to 8 pairs, spreading and curving upwards, not prominent on either surface ; length 1·75 to 4 in., breadth ·8 to 1·6 in., petiole ·25 to ·4 in. *Panicles* axillary and terminal, hoary-pubescent, many-flowered ; the flowers secund, 7 to 9 on each lateral branchlet, pedicelled, ·15 to ·2 in. long. *Sepals* sub-equal, broadly ovate, sub-acute, yellowish-pulverulent, tomentose externally, glabrous internally. *Petals* oblong, obtuse, twisted and with the apices reflexed in æstivation, spreading

when expanded, minutely yellowish-pulverulent, tomentose outside, glabrous inside. *Stamens* 15, sub-equal: apical awn curved, longer than the anther. *Ovary* ovoid, narrowing upwards into the style; stigma minute. *Fruit* ovoid, very slightly apiculate, striate, pale pubescent, about $\cdot 6$ in. long and $\cdot 3$ in. in diam., the persistent calyx covering the lower third of the fruit, sub-glabrous, thickened and concave at the base; the teeth deltoid, spreading. *Richetia penangiana*, Heim in Bull. Soc. Linn. Paris, 1891, p. 980.

Penang: on Government Hill, at an elevation of about 1,000 feet, Curtis, Nos. 1429 and 1393; Hullett, No. 188; King's Collector, No. 1534. Perak: King's Collector, Nos. 3333, 3707.

The leaves of this species, although larger, resemble those of *B. Curtisii*: but the fruits of the two are quite different. One of Mr. Curtis' specimens, No. 429 (communicated from Kew), forms the type of a new genus called *Richetia*, which M. Heim has founded (l. c. p. 975, also in his "Recherches sur les Dipterocarpaceés" p. 50), without having seen its flowers. I have retained for this M. Heim's specific name, while referring it to Beddome's older genus. The vernacular name of the species is *Dammar Etam*.

3. *BALANOCARPUS ANOMALUS*, King. A tree: young branches slender, dark-coloured, minutely lenticellate, the tips puberulous. *Leaves* coriaceous, ovate, acuminate; the base broad, sub-cuneate; both surfaces glabrous; main nerves 6 or 7 pairs, ascending, curving, not prominent: length 2·25 to 2·5 in., breadth 1 to 1·3 in., petiole $\cdot 6$ to $\cdot 7$ in. *Panicles* numerous, axillary and terminal, longer than the leaves, pubescent, their lateral branchlets bearing 6 to 8 sub-secund flowers. *Flowers* shortly pedicelled, $\cdot 15$ in. long. *Sepals* broadly ovate, connate at the base, obtuse, minutely tomentose outside, glabrous inside. *Petals* elliptic, blunt, yellowish adpressed-sericeous outside, glabrous inside, only about twice as long as the sepals, spreading and reflexed so as to expose the stamens and pistil. *Stamens* 10, in two rows; the filaments longer than the anthers, dilated; anther short, ovate, its connective produced into an apical awn as long as itself. *Ovary* ovoid, striate, pubescent, style short and thick, stigma small.

Kedah: Curtis, No. 1654.

Mr. Curtis is as yet the only collector of this, and his specimens have no fruit. I refer it to this genus, although its flowers differ from those of the other species known to me, in having petals only about twice as long as the sepals, spreading and reflexed so that the androgynœcium is quite exposed; and in having only 10 stamens. In other respects the specimens agree with *Balanocarpus*. Its vernacular name in Kedah is *Malaut*.

4. *BALANOCARPUS MAXIMUS*, King, n. sp. A tree 60 to 80 feet high : all parts except the inflorescence glabrous : young branches rather stout; the bark, loose, papery, lenticellate, pale. *Leaves* thinly coriaceous, oblong to elliptic-oblong, sub-acute, slightly narrowed to the rounded base; main nerves 7 to 9 pairs, slightly prominent beneath, the transverse veins slightly prominent when dry : length 5 to 7 in., breadth 2 to 2·5 in., petiole ·5 to ·6 in. *Panicles* axillary or terminal, about half as long as the leaves, few-flowered, minutely tomentose. *Flowers* subsessile, ·6 or ·7 in. long. *Sepals* broadly ovate, the outer two tomentose, the inner three more or less glabrous externally, all glabrous internally, the inner two with ciliate margins. *Petals* much longer than the sepals, narrowly oblong, the apex erose, expanded and concave at the base, adpressed-pubescent outside and towards the apex inside, otherwise glabrous. *Stamens* 10, in two rows; anthers with a deflexed terminal appendage from the connective. *Ovary* elongate, narrowly conical, sericeous. *Style* rather short, glabrous; stigma small. *Ripe fruit* cylindrical, tapering to each end but most to the apiculate apex; pericarp woody, striate, sub-glabrous, pale-brown when dry : 1·75 to 2·25 in. long, and ·6 or ·7 in. in diam. *Persistent sepals* fibrous, forming a toothed cup about ·5 in. deep, embracing the base of the fruit.

Perak : King's Collector, Nos. 7987 and 8006.

The flowers of this fine species do not exactly answer to Beddome's diagnosis of the genus *Balanocarpus*, inasmuch as they have 10 instead of 15 stamens, and neither of the cotyledons is lobed. In other respects the flowers and fruit agree perfectly.

5. *BALANOCARPUS HEIMII*, King n. sp. A tree 50 to 60 feet high : young branches rather slender, the bark dark-coloured, puberulous or glabrescent. *Leaves* coriaceous, narrowly oblong, tapering to the acuminate apex, and slightly narrowed to the rounded base; upper surface glabrous, shining, the midrib minutely pubescent: lower surface glabrescent except the pubescent midrib and 9 or 10 pairs of ascending, bold, slightly-curving nerves: length 4 to 6 in., breadth 1 to 1·75 in.; petiole ·3 or ·4 in., with minute black tomentum. *Flowers* unknown. *Ripe carpels* cylindrical, tapering to the apex, slightly narrowed to the base, 1·5 in. long and ·5 in. in diam.; the pericarp woody, sub-glabrous, sub-striate, dark-coloured when dry. *Persistent sepals* sub-equal, puberulous, thickened, forming a 5-lobed cup ·6 in. deep which embraces the base of the fruit. *Pierrea Penangiana*, Heim, MSS.

Penang : Curtis No. 273 (leaves only). Perak : King's Collector, No. 3718.

This tree, of which as yet only fruiting specimens have been found, so closely resembles the other Malayan species of *Balanocarpus* des-

cribed here, that I refer it without any hesitation to this genus. M. Curtis' leaf specimens of this have, I understand, received from M. Heim the MSS. name, *Pierrea penangiana*. The genus *Pierrea* has been founded by M. Heim (Bull. Soc. Linn. Paris, 1891, p. 958, and "Recherches sur les Diptercarpacées", p. 78) on specimens of which the author has not (as he admits) had the advantage of seeing the flowers. The vernacular name of this tree in Penang is *Chengah*, and its timber is, according to Mr. Curtis, very valuable. In the State of Perak, on the mainland almost opposite Penang, another species (*B. Wrayi*) receives a similar vernacular name.

6. *BALANOCARPUS WRAYI*, King n. sp. A tree: young branches slender, dark-coloured, glabrous. *Leaves* coriaceous, narrowly oblong, gradually tapering from the middle to the acute apex; the base sub-cuneate or rounded, slightly unequal-sided: both surfaces glabrous; main nerves 7 or 8 pairs, curved, oblique, slightly prominent beneath: length 2·25 to 2·75 in., breadth ·75 in.; petiole ·25 in., transversely wrinkled. *Panicles* axillary and terminal, nearly as long as the leaves. *Flowers* unknown. *Fruit* ovoid, much apiculate, glabrous, ·6 in. long, covered in its lower two-thirds by the persistent sub-acrescent glabrous calyx; outer two sepals smaller than the others, elliptic, obtuse, the inner three rotund, all thickened and concave.

Perak: Wray, No. 813.

Collected only once and without flowers. According to Mr. Wray the timber of this tree is valuable, and its vernacular name is *Chingi*, or *Chingal*. I refer this (in spite of the absence of flowers) to *Balanocarpus*, the other species of which it so closely resembles.

7. *BALANOCARPUS HEMSLEYANUS*, King, n. sp. A tree 50 to 100 feet high: young branches rather stout, rough, minutely lenticellate, puberulous. *Leaves* coriaceous, elliptic-oblong, sometimes slightly obovate, shortly and abruptly acuminate, slightly narrowed to the rounded or sub-emarginate base: upper surface glabrous except the minutely tomentose midrib; the lower scabrid from minute rigid stellately hairy tubercles which are most abundant on the stout midrib and nerves: main nerves 18 to 20 pairs, oblique, parallel, very prominent on the lower, obsolete on the upper, surface; length 7 to 12 in., breadth 3·25 to 5 in.; petiole ·6 to ·9 in. scabrid, pubescent. *Panicles* axillary or terminal, 3 to 7 in. long, scurfy stellate-pubescent; flowers rather crowded on the lateral branchlets, ·5 in. long, *Sepals* sub-equal, broadly ovate, acute, yellowish-tomentose externally, glabrous internally. *Petals* twice as long as the sepals, or longer, elliptic, oblique, obtuse, glabrous except a broad adpressed-sericeous band externally. *Stamens* 15, in three rows: the filaments dilated, unequal, longer than

the shortly ovate anthers; apical connectival appendage deflexed, curved, longer than the anther. *Ovary* elongated-conic, tomentose, tapering into the sparsely puberulous style; stigma small. *Ripe fruit* narrowly ovoid, apiculate, pale brownish-tomentose, 1.25 to 1.5 in. long. and .75 to 1 in. in diam. *Persistent sepals* nearly equal, their bases thickened, woody, pubescent, and concave, forming an irregularly 5-toothed cup which embraces the lower half of the fruit. *Shorea Hemsleyana*, King MSS. in Herb. Cal.

Penang: Curtis No. 2512. Perak: King's Collector, Nos. 5431, 6670, and 7562. Scortechini No. 1653.

This is an altogether anomalous species. It has leaves like several of the scabrid species of *Shorea*, such as *S. eximia* and *S. leprosula*. Its flowers are also more like those of *Shorea* than *Balanocarpus*; but its fruit is essentially that of the latter genus, in which, not without hesitation, I include it.

10. PACHYNOCARPUS, Hook. fil.

Resinous trees with the leaves and flowers of *Vatica*, but with sometimes only ten stamens. *Fruit* ovoid-globose, umbonate at the apex, 1-celled, 1-seeded, the pericarp densely coriaceous, splitting vertically. *Calyx* with five equal segments, at first almost free from the fruit, but the tube gradually accrescent, much thickened and adnate to the fruit, and finally embracing the whole of it except the apex. *Seed* pendulous, testa thin and adherent to the endocarp, cotyledons very thick and fleshy.

Leaves elliptic to oblong-elliptic, sub-acute or shortly and obtusely acuminate... .. 1. *P. Wallichii*.

Leaves broadly-elliptic or obovate-elliptic, the apex very blunt 2. *P. Stapfianus*.

Dr. Burck (in Ann. Jard. Bot. Buitenzorg) expands the definition of the genus *Vatica* so as to include not only the closely allied *Synaptea*, but also the genera *Isauxis* W. A., *Retinodendron*, Korth., and *Pachynocarpus* Hook. fil. To the union of *Synaptea* with *Vatica* I see no objection; for the whole difference between the two (as I have stated in a note under *Vatica*) consists in perfect freedom of the fruit in *Vatica* from the enlarged calyx, whereas in *Synaptea* there is a slight adhesion to the calyx at the very base. But for the inclusion of *Pachynocarpus*, I see no sufficient justification; for in this genus the calyx does not expand into membranous wings, but forms a dense fibro-cartilaginous cover for the fruit, which it tightly embraces, and to which it is quite adnate. As regards *Isauxis* and *Retinodendron*, they appear to me to be undistinguishable from each other by any but trivial marks, but they differ sufficiently in calyx from *Vatica* to be treated as a genus under the older name *Retinodendron*.

1. PACHYNOCARPUS WALLICHII, King. A tree 40 to 70 feet high: young branches deciduously scurfy-puberulous, their bark pale-brown, sparsely lenticellate. *Leaves* coriaceous, elliptic to oblong-elliptic,

sub-acute, or shortly and obtusely acuminate, the base cuneate; both surfaces glabrous, the lower pale and prominently reticulate when dry: main nerves 6 to 9 pairs, slightly prominent beneath, ascending; length 4·5 to 8 in., breadth 1·5 to 3 in., petiole; 4 to 6 in. *Panicles* crowded near the apices of the branches, many-flowered, 2 to 4 in. long. *Calyx-lobes* deltoid, minutely velvety outside. *Petals* linear-oblong, obtuse, puberulous externally. *Stamens* broadly ovoid, minutely but obtusely apiculate. *Ovary* puberulous: stigma sub-capitate, lobed. *Ripe fruit* ovoid-globose, about 75 in. in diam., closely embraced by the slightly shorter, much thickened, persistent, fibrous or woody, rugose, enlarged calyx-lobes. *V. Wallichii* Dyer in Journ. Bot. 1878 p. 154. *Vatica ruminata*, Burck in Ann. Jard. Bot. Buitenzorg, VI, 227 t. 29, fig. 4.

Penang: Wallich, Cat. No. 9018; Curtis Nos. 1161, 1218, 1391. Malacca: Maingay No. 201. Trang, King's Collector. Johore, Hullett and King. Perak: common at low elevations, King's Collector, Scortechini. **DISTRIB.**, Bangka.

In the young stages of the fruit of this species the calyx is quite small and embraces only the very base of it, much as in *Isauvis*; but as the fruit expands the calyx grows, so that when ripe the fruit is, with the exception of its apex, closely embraced by the much thickened, lignified, obscurely toothed calyx-tube. This offers, therefore, a transition between *Isauvis* and *Pachynocarpus*. And, indeed, it is to the former section that Dyer refers it (Journ. Bot., l. c.), and to which Burck refers his *D. ruminata*, a species which authentic specimens shew to be identical with this. Dr. Burck's species, *Vatica verrucosa* (Ann. Jard. Bot. Buitenzorg) appears also to come very near to this.

2. **PACHYNOCARPUS STAFFIANUS**, King, n. sp. A tree 80 to 100 feet high: young branches rather stout, scaly-pubescent at first, ultimately glabrous. *Leaves* coriaceous, broadly elliptic or obovate-elliptic, the apex broadly rounded, slightly narrowed to the rounded or sub-cuneate base: upper surface glabrous, shining, the lower paler, minutely and sparsely scurfy-puberulous on the midrib and nerves; main nerves 10 to 13 pairs, oblique, prominent on the lower, depressed on the upper, surface; length 5 to 8 in., breadth 2·75 to 4·5 in., petiole 65 to 1 in. *Flowers* unknown. *Ripe fruit* almost solitary, 2·5 to 3 in. long, on a woody raceme, globular, slightly apiculate, 1·25 in diam., closely invested by the gamosepalous, 5-toothed, thickened, woody, rugose, glaberulous calyx.

Perak: King's Collector, Nos. 5932 and 6132,

This very distinct species was first recognised as a *Pachynocarpus* by Dr. O. Stapf, of the Kew Herbarium, after whom I have named it. Its flowers are as yet unknown; but it is readily identified by its leaves.

11. ANCISTROCLADUS, Wall.

Smooth climbing shrubs with short supra-axillary, often arrested and circinate-hooked, branches. *Leaves* usually in terminal tufts, coriaceous, entire, reticulately feather-veined; exstipulate. *Flowers* usually small, very caducous, in terminal or lateral panicles. *Calyx-tube* at first short, adnate to the base of the ovary, its lobes imbricate, finally turbinate and adnate to the fruit, with the lobes unequally enlarged, spreading and membranous. *Stamens* 5 or 10, subperigynous. *Ovary* 1-celled, inferior; style sub-globose, persistent; *Stigmas* 3, erect, compressed, truncate, deciduous. Ovule solitary, erect or laterally affixed. *Seed* sub-globose, testa prolonged into the ruminations of the copious fleshy albumen; embryo short, straight; cotyledons short, divergent.—**DISTRIB.** Except *A. guineensis* in W. Tropical Africa, confined to Tropical Asia and the Indian Archipelago. Species about 10.

I follow the authors of the Genera Plantarum and the Flora of British India in including *Ancistrocladus* in *Dipterocarpeæ*. I venture, however, to think that it would be better to keep it as the type of a distinct Natural Order as MM. Planchon and De Candolle have done: for its characters do not fit well into the diagnosis of any other Order

1. ANCISTROCLADUS EXTENSUS, Wall. Cat. 1052. *Leaves* obovate or obovate-oblong, blunt or sub-acute, much narrowed at the base; panicles dichotomous, about half as long as the leaves: fruit smooth or slightly 5-ridged; accrescent calyx-lobes oblanceolate, obtuse. Planch. in Ann. Sc. Nat. Ser. 3, XIII, 318. DC. Prodr. XVI, 2, 602; Dyer in Hook. fil. Fl. Br. Ind. I, 299. *Ancistrolobus* sp. Griff. Notul. IV, 568, t. 605. fig. 2.

Andaman Islands. **DISTRIB.** Burmah.

Var. *pinangianus*; leaves sometimes oblanceolate-oblong, acute or sub-acuminate; panicles slender, lax, about as long as the leaves. *Ancistrocladus pinangianus*, Wall. Cat. 1054. Planchon in Ann. Sc. Nat. Ser. 3, XIII, 318; A. DC. Prodr. XVI. 2, 603; Dyer in Hook. fil. Fl. Br. Ind. I, 300.

Penang: Porter. Malacca: Maingay. (Kew Distrib.) No. 200. Singapore and Perak: King's Collectors. **DISTRIB.** Bangka, Sumatra, Burmah.